arXiv:math/0410017v1 [math.RT] 1 Oct 2004

REPRESENTATION THEORY OF THE DRINFEL'D DOUBLES OF A FAMILY OF HOPF ALGEBRAS

K. Erdmann E.L. Green* N. Snashall R. Taillefer†

Abstract

We investigate the Drinfel'd doubles $\mathcal{D}(\Lambda_{n,d})$ of a certain family of Hopf algebras. We determine their simple modules and their indecomposable projective modules, and we obtain a presentation by quiver and relations of these Drinfel'd doubles, from which we deduce properties of their representations, including the Auslander-Reiten quivers of the $\mathcal{D}(\Lambda_{n,d})$. We then determine decompositions of the tensor products of most of the representations described, and in particular give a complete description of the tensor product of two simple modules. This study also leads to explicit examples of Hopf bimodules over the original Hopf algebras.

Mathematics Subject Classification (2000): 17B37, 06B15, 81R50, 16W30, 16W35, 16G20, 16G70, 18D10.

1 Introduction

We study the representation theory of some Drinfel'd doubles. The Drinfel'd double of a finite-dimensional Hopf algebra was defined by Drinfel'd in order to provide solutions to the quantum Yang-Baxter equation arising from statistical mechanics.

Representations of a Hopf algebra (up to isomorphism) form a ring in which the product is given by the tensor product over the base field, and in the case of the Drinfel'd double (and any quasitriangular Hopf algebra) this ring is commutative.

However, not very much is known about the representations of the Drinfel'd double of a non-semisimple Hopf algebra in general. S. Witherspoon studied the Drinfel'd double of the algebra of a finite group in positive characteristic [W]. She proved in particular that the Green ring of the Drinfel'd double of a group algebra decomposes as a product of ideals associated to some subgroups of the original group. H-X. Chen gives a complete list of simple modules over the Drinfel'd doubles of the Taft algebras in [Ch]. D.E. Radford in [Rd2] studies the simple modules over the Drinfel'd double of a finite-dimensional Hopf algebra, and characterises them in some cases (which include the Taft algebras and quiver Hopf algebras); he then uses them to construct twist oriented quantum algebras, which give rise to invariants of oriented knots and links. Moreover, he establishes that all modules over the Drinfel'd double can be viewed as submodules of some particular Yetter-Drinfel'd modules (constructed from modules over the original Hopf algebra).

An important property of the algebra underlying the Drinfel'd double of a finite-dimensional Hopf algebra H can be deduced from work of D.E. Radford and R. Farnsteiner ([Rd, Corollary 2 and Theorem 4] and [F, Proposition 2.3]): the Drinfel'd double $\mathcal{D}(H)$ is a symmetric algebra.

In this paper, we study in detail the representation theory of the Drinfel'd doubles of the algebras $\Lambda_{n,d}$, that is, of the duals of the extended Taft algebras (defined in the first section). The algebras $\Lambda_{n,d}$ are interesting, since they are finite-dimensional Hopf algebras which are neither commutative nor cocommutative, and which are not quasitriangular, with antipodes of arbitrarily high order ($2n$). However, they are monomial (that is, all the relations are given by paths) and have finite representation type, and as such are therefore very suitable to study.

In the first section, we study the representations of these Drinfel'd doubles, first describing the projective modules and the simple modules (see 2.15, 2.16, 2.20 and 2.24). We give a complete presentation by quiver and relations, and this shows that the algebras are of tame representation

---

*Partially supported by an NSA grant

†Partially supported by an LMS Scheme 4 grant and a Lavoisier grant from the French Ministry of Foreign Affairs

type (see Section 3; see also [B, 4.4] for definitions of representation types). These quivers are similar to some quivers which occur in the representation theory of blocks of reduced enveloping algebras in characteristic p (see [FS]). However, the situation in our case is quite different: as we see later on in the paper, there exist representations which are periodic of period greater than 2 in general (which is not the case for reduced enveloping algebras). Moreover, the conditions on the characteristic of the field in our context and in the case of reduced enveloping algebras are distinct; in the case of reduced enveloping algebras, the cycles which occur in the quiver have a length which is a power of the characteristic, whereas in our case the characteristic does not divide the length of the cycles but is otherwise arbitrary.

In the following section we describe all the indecomposable representations of $\mathcal{D}(\Lambda_{n,d})$, and the Auslander-Reiten quiver of these algebras. We also determine which of these representations are splitting trace modules and more precisely give the quantum dimensions of the modules (see Section 3.3, where these concepts are also defined). We then study tensor products of representations. We first give a complete and explicit decomposition of the tensor product of two simple modules, and in particular prove that this is always semisimple modulo some projective direct summands (see 4.1). We then describe the tensor products of many other modules up to projectives. This gives a large part of the structure of the Green ring of $\mathcal{D}(\Lambda_{n,d})$ modulo projectives. Finally, we give some examples of Hopf bimodules over the original Hopf algebras $\Lambda_{n,d}$, obtained via the equivalence of categories between Hopf bimodules over $\Lambda_{n,d}$ and modules over $\mathcal{D}(\Lambda_{n,d})$. We conclude with an appendix in which we outline the proof of the classification of the representations of $\mathcal{D}(\Lambda_{n,d})$.

This work opens many questions on the representation theory of Drinfel'd doubles, in particular the possibility of finding some general properties of tensor products of simple modules over the Drinfel'd doubles of some Hopf algebras, using D.E. Radford's results [Rd2].

For general facts about representations of algebras, we refer to [ARS], [B] and [E]. We will refer to those more precisely in some parts of the paper.

Acknowledgements: The second and third authors thank the University of Oxford and the last author thanks the Virginia Polytechnic Institute and State University for their kind hospitality.

2 Projectives and quiver for $\mathcal{D}(\Lambda_{n,d})$

2.1 Preliminaries

Throughout the paper, k is an algebraically closed field.

The algebra $\Lambda_{n,d}$ is described by quiver and relations; we refer to [ARS, III.1] and [B, 4.1] for definitions and properties relating to quivers and relations. The quiver is cyclic,



with n vertices $e_0, \ldots, e_{n-1}$ and n arrows $a_0, \ldots, a_{n-1}$, where the arrow a_i goes from the vertex e_i to the vertex e_{i+1}, and we factor by the ideal generated by all paths of length $d \geqslant 2$. We shall denote by γ_i^m the path $a_{i+m-1} \cdots a_{i+1} a_i$ (read from right to left), that is, the path of length m starting at the vertex e_i.

When d divides n, this algebra is a Hopf algebra, and in fact the condition $d \mid n$ is a necessary and sufficient condition for $\Lambda_{n,d}$ to be a Hopf algebra when $\mathrm{char} k = 0$ (see [Ci, CHZ]). This Hopf algebra can actually be considered over more general fields, and in this paper, we assume only that the characteristic of k does not divide n (to ensure existence of roots of unity).

We fix a primitive d^{th} root of unity q in k. The formulae

$$\begin{array}{lll} \varepsilon(e_i) = \delta_{i0} & \Delta(e_i) = \sum_{j+\ell=i} e_j \otimes e_\ell & S(e_i) = e_{-i} \\ \varepsilon(a_i) = 0 & \Delta(a_i) = \sum_{j+\ell=i}(e_j \otimes a_\ell + q^\ell a_j \otimes e_\ell) & S(a_i) = -q^{i+1}a_{-i-1} \end{array}$$

determine the Hopf algebra structure of $\Lambda_{n,d}$.

We want to study the Drinfel'd double of $\Lambda_{n,d}$. Recall:

Definition 2.1 (see [Kl, M]) *Let H be a Hopf algebra. The* **Drinfel'd double** *of H is the Hopf algebra which is equal to $H^{*cop} \otimes H$ as a coalgebra (ordinary tensor product of coalgebras), and whose product is defined by*

$$(\alpha \otimes h)(\beta \otimes g) = \alpha\beta(S^{-1}h^{(3)}?h^{(1)}) \otimes h^{(2)}g,$$

where $\beta(S^{-1}h^{(3)}?h^{(1)})$ is the map which sends $x \in H$ to $\beta(S^{-1}h^{(3)}xh^{(1)}) \in k$ and where we have used the Sweedler notation $\Delta(h) = h^{(1)} \otimes h^{(2)}$ for the comultiplication.

Therefore, we need to understand the dual $\Lambda_{n,d}^{*cop}$; it is isomorphic as an algebra to the extended Taft algebra

$$\langle G, X \mid G^n = 1, X^d = 0, GX = q^{-1}XG\rangle.$$

(C. Cibils explains this for $n = d$ when $\Lambda_{n,d}$ is a selfdual Hopf algebra in [Ci], and the general case is similar; if $\{\check{\gamma}_i^m \mid i \in \mathbb{Z}_n, 0 \leqslant m \leqslant d-1\}$ denotes the dual basis of $\Lambda_{n,d}^{*cop}$, the correspondence is determined by $\check{e}_i \mapsto G^i$ and $\check{a}_i \mapsto G^iX$). Its Hopf algebra structure is defined by

$$\begin{array}{lll} \varepsilon(G) = 1 & \Delta(G) = G \otimes G & S(G) = G^{-1} \\ \varepsilon(X) = 0 & \Delta(X) = X \otimes G + 1 \otimes X & S(X) = -XG^{-1} = -q^{-1}G^{-1}X. \end{array}$$

Notation **2.2** We shall need the following notation to describe some elements in $\mathcal{D}(\Lambda_{n,d})$: for non-negative integers m and u, we define the **q-integers** $(0)_q = 0$ and $(m)_q = 1 + q + q^2 + \ldots + q^{m-1}$, the **$q$-factorials** $0!_q = 1$ and $m!_q = (m)_q(m-1)_q \ldots (2)_q(1)_q$, and the **$q$-binomial coefficients** $\binom{m}{u}_q = \dfrac{m!_q}{(m-u)!_q u!_q}$ (see for instance [Ci, Rd3], where some properties are also given).

Remark **2.3** Using the notation above, we have $\Delta(\gamma_i^m) = \sum\limits_{\substack{j \in \mathbb{Z}_n \\ 0 \leqslant v \leqslant m}} \binom{m}{v}_q q^{vj}\gamma_{i-j}^v \otimes \gamma_j^{m-v}$ and $\Delta(X^m) = \sum_{v=0}^m \binom{m}{v}_q X^v \otimes G^vX^{m-v}$.

We can now describe the Drinfel'd double $\mathcal{D}(\Lambda_{n,d})$:

Proposition 2.4 *The Drinfel'd double $\mathcal{D}(\Lambda_{n,d})$ is described as follows: as a coalgebra, it is $\Lambda_{n,d}^{*cop} \otimes \Lambda_{n,d}$. We write the basis elements $G^iX^j\gamma_\ell^m$, with $i, \ell \in \mathbb{Z}_n$ and $0 \leqslant j, m \leqslant d-1$ (i.e. we do not write the tensor product symbol). The following relations determine the algebra structure completely:*

$G^n = 1, X^d = 0,$ *and* $GX = q^{-1}XG$,

the product of elements γ_ℓ^m is the usual product of paths,

$\gamma_\ell^m G = q^{-m}G\gamma_\ell^m$, *and*

$\gamma_\ell^m X = q^{-m}X\gamma_{\ell+1}^m - q^{-m}(m)_q\gamma_{\ell+1}^{m-1} + q^{\ell+1-m}(m)_q G\gamma_{\ell+1}^{m-1}$.

Proof: The proof is straightforward, using the definition of the Drinfel'd double (Definition 2.1) and the structure of $\Lambda_{n,d}^{*cop}$. ■

Notation **2.5** Since some indices are described modulo d and others modulo n, we need to make a distinction. If j is an element in $\mathbb{Z}_n$, we shall denote its representative modulo d in $\{1, \ldots, d\}$ by $\langle j\rangle$ and its representative modulo d in $\{0, \ldots, d-1\}$ by $\langle j\rangle^-$.

2.2 Quiver of $\mathcal{D}(\Lambda_{n,d})$

The aim of this section is to describe the quiver of $\mathcal{D}(\Lambda_{n,d})$. To do this, we first decompose $\mathcal{D}(\Lambda_{n,d})$ into a (non-minimal) product of algebras $\Gamma_0, \ldots, \Gamma_{n-1}$, and study each of these algebras. We give a basis for Γ_u for $u = 0, \ldots, n-1$, and describe some indecomposable Γ_u-modules as a first step in the construction of the indecomposable projective modules. This follows R. Suter's method in [S] where he studies the representations of a finite-dimensional quotient of $U_q(\mathfrak{sl}_2(\mathbb{C}))$ (see also [X, P]).

2.2.1 First decomposition of $\mathcal{D}(\Lambda_{n,d})$

Proposition 2.6 *The elements* $E_u := \frac{1}{n}\sum_{i,j\in\mathbb{Z}_n} q^{-i(u+j)}G^i e_j$, *for* $u \in \mathbb{Z}_n$, *are central orthogonal idempotents, and* $\sum_{u\in\mathbb{Z}_n} E_u = 1$.

Therefore $\mathcal{D}(\Lambda_{n,d}) \cong \prod_{u\in\mathbb{Z}_n}\Gamma_u$ *where* $\Gamma_u = \mathcal{D}(\Lambda_{n,d})E_u$.

We shall now study Γ_u.

2.2.2 Construction of modules over $\mathcal{D}(\Lambda_{n,d})$

We now define some idempotents inside Γ_u, which are not central, but we will use them to describe a basis for Γ_u.

Proposition 2.7 *Set* $E_{u,j} = \sum_{v=0}^{\frac{n}{d}-1} e_{j+vd}E_u$, *for* $j \in \mathbb{Z}_d$. *Then* $E_{u,j}E_{u,\ell} = \delta_{j\ell}E_{u,j}$ *and* $\sum_{j=0}^{d-1} E_{u,j} = E_u$. *We also have* $E_{u,j} = E_{u,j'}$ *iff* $j \equiv j' \pmod{d}$.

Moreover, the following relations hold within Γ_u*:*

$$GE_{u,j} = q^{u+j}E_{u,j} = E_{u,j}G \qquad XE_{u,j} = E_{u,j-1}X$$

$$\gamma_\ell^m E_{u,j} = \begin{cases} E_{u,j+m}\gamma_\ell^m & \text{if } \ell \equiv j \pmod{d} \\ 0 & \text{otherwise.} \end{cases}$$

We can now describe a basis for Γ_u and a grading on Γ_u, as follows: $\Gamma_u = \bigoplus_{s=-d+1}^{d-1}(\Gamma_u)_s$ with $(\Gamma_u)_s = \operatorname{span}\{X^t\gamma_j^m E_{u,j} \mid j \in \mathbb{Z}_n, 0 \leqslant m, t \leqslant d-1, m-t = s\}$.

Multiplication on the left and on the right by elements in $\mathcal{D}(\Lambda_{n,d})$ respect this grading, with multiplication by X on either side reducing the degree by 1, multiplication by an arrow on either side increasing the degree by 1 and multiplication by G on either side leaving the degree unchanged. Moreover $(\Gamma_u)_s$ is a sum of eigenspaces for G : if $yE_{u,j}$ is an element in $(\Gamma_u)_s$, we have $G \cdot yE_{u,j} = q^{s+j+u}yE_{u,j}$ and $yE_{u,j} \cdot G = q^{j+u}yE_{u,j}$.

Now set $F_{u,j} := \gamma_j^{d-1}E_{u,j}$ for $j \in \mathbb{Z}_n$. Then:

Proposition 2.8 *The module* $\Gamma_u F_{u,j}$ *has the following form:*

$$\begin{tikzcd}
F_{u,j} \arrow[d, shift left, "X"] \\
\bullet \arrow[u, shift left, "a_{j+d-2}"] \arrow[d, shift left, "X"] \\
\bullet \arrow[u, shift left, "a_{j+d-3}"] \arrow[d, dotted, no head] \\
\bullet \arrow[d, shift left, "X"] \\
H_{u,j} \arrow[u, shift left, "a"] \arrow[d, "X"] \\
\tilde{F}_{u,j} \arrow[d, shift left, "X"] \\
\bullet \arrow[u, shift left, "a"] \arrow[d, dotted, no head] \\
\bullet \arrow[d, shift left, "X"] \\
\tilde{H}_{u,j} \arrow[u, shift left, "a_j"]
\end{tikzcd}$$

where $H_{u,j} := X^{\langle 2j+u-1\rangle -1}F_{u,j}$, $\tilde{F}_{u,j} := X^{\langle 2j+u-1\rangle^-}F_{u,j}$ *and* $\tilde{H}_{u,j} := X^{d-1}F_{u,j}$ *(recall the notations in 2.5).*

In this diagram, the arrows represent the actions of X *and of the arrows in the original quiver up to a nonzero scalar; the basis vectors are eigenvectors for the action of* G.

Note that when $2j+u-1 \equiv 0 \pmod{d}$, *the single arrow does not occur, the module is simple, and we have* $H_{u,j} = \tilde{H}_{u,j} = X^{d-1}F_{u,j}$, *and* $\tilde{F}_{u,j} = F_{u,j}$.

In order to prove this proposition, we require the following lemma:

Lemma 2.9 *The element* $\gamma^b_{d+j-m-1}X^mF_{u,j}$ *is equal to*

$$q^{-\frac{b(2m-b+1)}{2}}\frac{(m)!_q}{(m-b)!_q}\prod_{t=1}^{b}(q^{2j+u-1-(m-t+1)}-1)\,X^{m-b}F_{u,j}$$

if $b<m$ *and is 0 otherwise.*

Proof: This is proved by induction on b. When $b=1$, we take the arrow on the left across the X's, using the relations in Proposition 2.4.

$$\begin{aligned}
a_{j-m-1+d}X^m\gamma_j^{d-1}E_{u,j} &= \left(q^{-1}Xa_{j-m+d} - q^{-1}(1-q^{2(j-m+d)+u+2})e_{j-m+d}\right)X^{m-1}\gamma_j^{d-1}E_{u,j}\\
&= q^{-1}\left(q^{-1}X^2a_{j-m+d+1} - q^{-1}(1-q^{2(j-m+d)+u+2})X\right)X^{m-2}\gamma_j^{d-1}E_{u,j}\\
&\quad -q^{-1}(1-q^{2(j-m+d-1)+u+2})X^{m-1}\gamma_j^{d-1}E_{u,j}\\
&= q^{-2}X^2a_{j-m+d+1}X^{m-2}\gamma_j^{d-1}E_{u,j}\\
&\quad +\left(-q^{-2}-q^{-1}+q^{2j-2m+u-1}+q^{2j-2m+u}\right)X^{m-1}\gamma_j^{d-1}E_{u,j}\\
&= \dots\\
&= q^{-m}X^ma_{j+d-1}\gamma_j^{d-1}E_{u,j}\\
&\quad +\sum_{p=1}^{m}(-q^{-p}+q^{2j-2m+u+p-2})X^{m-1}\gamma_j^{d-1}E_{u,j},
\end{aligned}$$

with the first term in the last identity equal to 0. ■

Proof of Proposition **2.8:** Here we apply Lemma 2.9 with $b=1$ to see that the element $a_{d+j-m-1}X^mF_{u,j}$ is equal to 0 if $m \equiv 2j+u-1 \pmod{d}$ and is a nonzero multiple of $X^{m-1}F_{u,j}$ otherwise. ■

Remark **2.10** The simple projective modules that we have described in Proposition 2.8 when $2j+u-1 \equiv 0 \pmod{d}$, are distributed as follows in the whole algebra $\mathcal{D}(\Lambda_{n,d})$:

When d is odd, there are $\frac{n}{d}$ such simples in each algebra Γ_u.

When d is even, there are $\frac{2n}{d}$ such simples in Γ_u if u is odd, and none in Γ_u for u even.

Definition 2.11 *We define permutations of the indices in* $\mathbb{Z}_n$ *by* $\sigma_u(j) = d+j-\langle 2j+u-1\rangle$. *Note that the arrow going up from* $H_{u,j}$ *in the diagram in Proposition 2.8 is* $a_{\sigma_u(j)}$.

Remark **2.12** We can easily see that $\sigma_u(j) = j$ if and only if $2j+u-1 \equiv 0 \pmod{d}$, and that if $2j+u-1 \not\equiv 0 \pmod{d}$, then $\sigma_u^2(j) = j+d$ and so σ_u has order $\frac{2n}{d}$.

We shall now define larger modules (and we will see later that they are a full set of representatives of the indecomposable projective modules).

2.2.3 Indecomposable projective modules and decomposition of $\mathcal{D}(\Lambda_{n,d})$

Lemma 2.13 *If* $2j+u-1 \not\equiv 0 \pmod d$, *then there exists an element* $K_{u,j}$ *homogeneous of degree* $d-\langle 2j+u-1\rangle^- - 1$ *such that* $H_{u,j} = a_{\sigma_u(j)-1}K_{u,j}$.

Proof: Consider $H_{u,j} = X^{\langle 2j+u-1\rangle -1}\gamma_j^{d-1}E_{u,j}$. We first take one arrow across the X's; there exist scalars $\alpha_1,\dots,\alpha_{2j+u-2}$ in k such that:

$$\begin{aligned}
H_{u,j} &= X^{\langle 2j+u-1\rangle-1}\gamma_j^{d-1}E_{u,j}\\
&= qX^{\langle 2j+u-1\rangle-2}a_{j+d-1}X\gamma_j^{d-2}E_{u,j} + \alpha_1 X^{\langle 2j+u-1\rangle-2}\gamma_j^{d-2}E_{u,j}\\
&= q^2X^{\langle 2j+u-1\rangle-3}a_{j+d}X^2\gamma_j^{d-2}E_{u,j} + (\alpha_1+\alpha_2) X^{\langle 2j+u-1\rangle-2}\gamma_j^{d-2}E_{u,j}\\
&= \dots\\
&= q^{2j+u-2}a_{\sigma_u(j)-1}X^{\langle 2j+u-1\rangle-1}\gamma_j^{d-2}E_{u,j} + (\alpha_1+\dots+\alpha_{2j+u-2}) X^{\langle 2j+u-1\rangle-2}\gamma_j^{d-2}E_{u,j}.
\end{aligned}$$

We now repeat the process on the second term of the last identity, and continue until there is an arrow in front of all the terms; so there exist scalars β_i', β_i'' and β_i such that the following identities hold:

$$\begin{aligned}
H_{u,j} &= q^{2j+u-2}a_{\sigma_u(j)-1}X^{\langle 2j+u-1\rangle-1}\gamma_j^{d-2}E_{u,j} + \beta_1' X^{\langle 2j+u-1\rangle-2}\gamma_j^{d-2}E_{u,j}\\
&= q^{2j+u-2}a_{\sigma_u(j)-1}X^{\langle 2j+u-1\rangle-1}\gamma_j^{d-2}E_{u,j}\\
&\quad +\beta_1'\left(q^{2j+u-3}a_{\sigma_u(j)-1}X^{\langle 2j+u-1\rangle-2}\gamma_j^{d-3}E_{u,j} + \beta_2'' X^{\langle 2j+u-1\rangle-3}\gamma_j^{d-3}E_{u,j}\right)\\
&= q^{2j+u-2}a_{\sigma_u(j)-1}X^{\langle 2j+u-1\rangle-1}\gamma_j^{d-2}E_{u,j} + \beta_1 a_{\sigma_u(j)-1}X^{\langle 2j+u-1\rangle-2}\gamma_j^{d-3}E_{u,j}\\
&\quad +\beta_2' X^{\langle 2j+u-1\rangle-3}\gamma_j^{d-3}E_{u,j}\\
&= \dots\\
&= q^{2j+u-2}a_{\sigma_u(j)-1}X^{\langle 2j+u-1\rangle-1}\gamma_j^{d-2}E_{u,j} + a_{\sigma_u(j)-1}\sum_{p=2}^{\langle 2j+u-1\rangle}\beta_{p-1}X^{\langle 2j+u-1\rangle-p}\gamma_j^{d-p-1}E_{u,j}\\
&= a_{\sigma_u(j)-1}K_{u,j}
\end{aligned}$$

with $K_{u,j}$ nonzero and homogeneous of degree $d-\langle 2j+u-1\rangle - 1 = d-\langle 2j+u-1\rangle^- - 1$.

■

Definition 2.14 *If* $2j+u-1\equiv 0 \pmod d$, *set* $K_{u,j} = F_{u,j} (= \tilde{F}_{u,j})$. *Note that it is homogeneous of degree* $d-1$.

Now consider $\Gamma_u K_{u,j}$. The following result is immediate:

Definition-Proposition 2.15 *Assume that* $2j+u-1\equiv 0 \pmod d$. *Define* $L(u,j) := \Gamma_u K_{u,j}$; *this module has the following structure:*



In the case $2j+u-1\not\equiv 0 \pmod d$, we obtain the following structure:

Proposition 2.16 *Assume that $2j+u-1 \not\equiv 0 \pmod{d}$. The module $\Gamma_u K_{u,j}$ has the following structure:*



To prove this, we require the following two lemmas:

Lemma 2.17 *We have* $\gamma^{t}_{\sigma_u(j)-s-1} X^s K_{u,j} = \sum_{b=s-t+1}^{s} q^{-b} c_{b,s} \gamma^{b-s+t-1}_{\sigma_u(j)-b} X^b H_{u,j} + c_{s-t,s} X^{s-t} K_{u,j}$ *where* $c_{s,s}=1$, $c_{b,s}=0$ *if* $b<0$, *and* $c_{b,s}=\zeta_s\zeta_{s-1}\dots\zeta_{b+1}$ *with* $\zeta_s=(s)_q q^{-s}(q^{-(2j+u-1)-s}-1)$ *if* $0\leqslant b\leqslant s-1$.

Proof: The proof is by induction on t, and we write out the case $t=1$ here:

$$\begin{aligned}
a_{\sigma_u(j)-s-1} X^s K_{u,j} &= q^{-1} X a_{\sigma_u(j)-s} X^{s-1} K_{u,j} - q^{-1}(1-q^{2j-2\langle 2j+u-1\rangle-2s+u}) X^{s-1} K_{u,j} \\
&= q^{-1} X a_{\sigma_u(j)-s} X^{s-1} K_{u,j} - (q^{-1}-q^{-2j-u+1-2s}) X^{s-1} K_{u,j} \\
&= q^{-1}\left(q^{-1} X^2 a_{\sigma_u(j)-s+1} X^{s-2} K_{u,j} - (q^{-1}-q^{-2j-u+1-2s+2}) X^{s-1} K_{u,j}\right) \\
&\quad -(q^{-1}-q^{-2j-u+1-2s}) X^{s-1} K_{u,j} \\
&= q^{-2} X^2 a_{\sigma_u(j)-s+1} X^{s-2} K_{u,j} \\
&\quad -(q^{-1}+q^{-2}-q^{-2j-u+1-2s}-q^{-2j-u+1-2s+1}) X^{s-1} K_{u,j} \\
&= \dots \\
&= q^{-s} X^s a_j K_{u,j} + \zeta_s X^{s-1} K_{u,j}.
\end{aligned}$$

■

Lemma 2.18 *For* $0\leqslant s\leqslant d-\langle 2j+u-1\rangle^{-}-1$, *the elements* $X^s K_{u,j}$ *and* $X^s \tilde{F}_{u,j}$ *are linearly independent (for other values of* s, *we have* $X^s\tilde{F}_{u,j}=0$*).*

Proof: Assume that $\alpha X^s K_{u,j} + \beta X^s \tilde{F}_{u,j} = 0$ with $0 \leqslant s \leqslant d - \langle 2j+u-1\rangle^- - 1$. Multiply by $\gamma^{s+1}_{\sigma_u(j)-1-s}$ using Lemma 2.17 and Lemma 2.9: then $\gamma^{s+1}_{\sigma_u(j)-1-s} X^s \tilde{F}_{u,j}$ is a multiple of $(q^{-2j+s+\langle 2j+u-1\rangle-u+2-(s+1)} - 1)X^{\langle 2j+u-1\rangle} F_{u,j}$ which is zero.

On the other hand, $\gamma^{s+1}_{\sigma_u(j)-1-s} X^s K_{u,j}$ is equal to $\sum_{b=0}^{s} q^{-b} c_{b,s} \gamma^{b}_{\sigma_u(j)-b} X^b H_{u,j}$. Now $\zeta_p = 0$ if and only if $p \equiv -2j-u+1$; but if $0 \leqslant b+1 \leqslant p \leqslant s \leqslant d - \langle 2j+u-1\rangle^- - 1$, we have $\zeta_p \neq 0$ and therefore $c_{b,s} \neq 0$ for all b, s with $0 \leqslant b+1 \leqslant s \leqslant d - \langle 2j+u-1\rangle^- - 1$.

So multiplying the identity $\alpha X^s K_{u,j} + \beta X^s \tilde{F}_{u,j} = 0$ by $\gamma^{s+1}_{\sigma_u(j)-1-s}$ gives a nonzero multiple of $\alpha \gamma^{s+1}_{\sigma_u(j)-1-s} X^s K_{u,j}$ with $\gamma^{s+1}_{\sigma_u(j)-1-s} X^s K_{u,j}$ nonzero, so $\alpha = 0$ and therefore $\beta = 0$. ■

Proof of Proposition **2.16:** We apply Lemma 2.17 with $t = d-1 = s$ to see that $\gamma^{d-1}_{\sigma_u(j)-d} X^{d-1} K_{u,j}$ is a nonzero multiple of $\tilde{F}_{u,j}$: if $b \leqslant d - \langle 2j+u-1\rangle^- - 1$ then $c_{b,d-1} = 0$; if $b \geqslant d - \langle 2j+u-1\rangle^- + 1$ then $\gamma^{b-1}_{\sigma_u(j)-b} X^b H_{u,j} = 0$; finally, if $b = d - \langle 2j+u-1\rangle^-$, then $\gamma^{d-1}_{\sigma_u(j)-d} X^{d-1} K_{u,j}$ is a nonzero multiple of $X^{\langle 2j+u-1\rangle} \gamma^{d-1}_{j} E_{u,j}$. In particular, $X^s K_{u,j} \neq 0$ for all $s = 0, \ldots, d-1$. The rest of the structure follows from this and Lemma 2.18. ■

We can easily find all the submodules of $\Gamma_u K_{u,j}$, and therefore:

Definition-Proposition 2.19 *When* $2j+u-1 \not\equiv 0 \pmod d$, *the module* $\Gamma_u K_{u,j}$ *has exactly two composition series:*

$$\Gamma_u K_{u,j} \supset \Gamma_u F_{u,j} + \Gamma_u \tilde{D}_{u,j} \supset \Gamma_u F_{u,j} \supset \Gamma_u \tilde{F}_{u,j} \supset 0$$

and

$$\Gamma_u K_{u,j} \supset \Gamma_u F_{u,j} + \Gamma_u \tilde{D}_{u,j} \supset \Gamma_u \tilde{D}_{u,j} \supset \Gamma_u \tilde{F}_{u,j} \supset 0.$$

Define $L(u,j) := \frac{\Gamma_u K_{u,j}}{\Gamma_u F_{u,j} + \Gamma_u \tilde{D}_{u,j}}$; *this is a simple module of dimension* $d - \langle 2j+u-1\rangle^-$, *and the composition factors of the composition series above are* $L(u,j)$, $L(u,\sigma_u(j))$, $L(u,\sigma_u^{-1}(j))$, *and* $L(u,j)$.

Proof: The proof is straightforward using Proposition 2.16 and the Proposition 2.20 which follows. ■

Proposition 2.20 *Let* S *be a simple module. Then* S *is isomorphic to* $L(u,j)$ *if and only if the three following properties hold:*

(a) Dimensions: $\dim S = \dim L(u,j)$.

(b) Action of the central idempotents: E_u *acts as identity on* S, *and* E_v *acts as zero on* S *if* $v \neq u$.

(c) Let Y *be the generator of* S *which is in the kernel of the action of* X *(this is well-defined up to a scalar and corresponds to* $\tilde{H}_{v,j}$*). Then the vertex* e_j *acts as identity on* Y, *and the other vertices act as zero.*

Note that the action of G *on this same element* Y *is multiplication by* q^{j+u}.

Moreover, $\dim L(u,\sigma_u^t(j)) = \begin{cases} d - \langle 2j+u-1\rangle^- & \text{if } t \text{ is even} \\ \langle 2j+u-1\rangle & \text{if } t \text{ is odd.} \end{cases}$

To summarize:

Proposition 2.21 *If* $2j+u-1 \not\equiv 0 \pmod d$, *then* $\Gamma_u K_{u,j} = \bigoplus_{h=0}^{d-1} \left(kX^h F_{u,j} \oplus kX^h K_{u,j}\right)$ *as a vector space, it has dimension* $2d$, *and it is an indecomposable* Γ_u*-module with simple top and simple socle, both isomorphic to* $L(u,j)$.

If $2j+u-1 \equiv 0 \pmod d$, *then* $L(u,j) := \Gamma_u K_{u,j} = \bigoplus_{h=0}^{d-1} kX^h F_{u,j}$ *as a vector space, it has dimension* d, *and it is a simple* Γ_u*-module.*

To decompose Γ_u into a sum of indecomposable modules, we find modules isomorphic to the $\Gamma_u K_{u,j}$ inside Γ_u :

Lemma 2.22 *If $0 \leqslant h \leqslant d - \langle 2j+u-1\rangle^{-} - 1$, then right multiplication by X^h induces an isomorphism $\Gamma_u K_{u,j} \xrightarrow{\sim} \Gamma_u K_{u,j} X^h$ of Γ_u-modules.*

Proof: We must show that right multiplication by X^h maps $\Gamma_u K_{u,j}$ injectively in $\Gamma_u K_{u,j} X^h$, and it is enough to do this for h maximal. For this, we only need to check that $\tilde{H}_{u,j} X^h \neq 0$ and $\tilde{D}_{u,j} X^h \neq 0$. Since $\tilde{H}_{u,j} X^h \neq 0$ implies $\tilde{D}_{u,j} X^h \neq 0$, and since $h \leqslant d - \langle 2j+u-1\rangle^{-} - 1$, we only need to consider $\tilde{H}_{u,j} X^{d-\langle 2j+u-1\rangle^{-}-1}$.

To compute this, we need a relation similar to that in Lemma 2.9, which is proved in the same way: we have

$$X^{d-1}\gamma_j^m E_{u,j} X^b = q^{-\frac{b(2m-b+1)}{2}} \frac{(m)!_q}{(m-b)!_q} \prod_{t=1}^{b} (q^{2j+m+u+t} - 1)\, X^{d-1}\gamma_{j+b}^{m-b} E_{u,j+b}$$

if $b < m$ and $X^{d-1}\gamma_j^m E_{u,j} X^b = 0$ otherwise. Using this relation, we see that $\tilde{H}_{u,j} X^{d-\langle 2j+u-1\rangle^{-}-1}$ is a nonzero multiple of $\prod_{t=1}^{d-\langle 2j+u-1\rangle^{-}-1} (q^{2j-1+u+t} - 1) X^{d-1} \gamma_{j+d-\langle 2j+u-1\rangle^{-}-1}^{\langle 2j+u-1\rangle^{-}} E_{u,-j-u}$ which is nonzero. ■

We can now decompose Γ_u entirely into a sum of indecomposable Γ_u-modules:

Theorem 2.23 *Γ_u decomposes into a direct sum of indecomposable modules in the following way:*

$$\Gamma_u = \bigoplus_{j \in \mathbb{Z}_n} \bigoplus_{h=0}^{d-\langle 2j+u-1\rangle^{-}-1} \Gamma_u K_{u,j} X^h.$$

Proof: We first prove that the sum is direct: the sums over h (for j fixed) are direct because the summands are in different right G-eigenspaces ($K_{u,j} X^h G = K_{u,j} X^h G E_{u,j+h} = q^{u+j+h} K_{u,j} X^h$). The outer sum is also direct because the summands have non-isomorphic socles: the socle of $\bigoplus_{h=0}^{d-\langle 2j+u-1\rangle^{-}-1} \Gamma_u K_{u,j} X^h$ is $\bigoplus_{h=0}^{d-\langle 2j+u-1\rangle^{-}-1} L(u,j) X^h$ with $L(u,j)X^h \cong L(u,j)$.

Equality follows from dimension counting. ■

Corollary 2.24 *Set $P(u,j) = \Gamma_u K_{u,j}$ for all u, j. The modules $P(u,j)$ are projective, and they represent the different isomorphism classes of projective $\mathcal{D}(\Lambda_{n,d})$-modules when u and j vary in $\mathbb{Z}_n$.*

When $2j + u - 1 \not\equiv 0 \pmod{d}$, their structure is

$$\begin{tikzcd}
 & L(u,j) \arrow[dl, dash] \arrow[dr, dash] & \\
L(u,\sigma_u^{-1}(j)) \arrow[dr, dash] & & L(u,\sigma_u(j)) \arrow[dl, dash] \\
 & L(u,j) &
\end{tikzcd}$$

and when $2j + u - 1 \equiv 0 \pmod{d}$, $P(u,j) = L(u,j)$ is simple of dimension d.

Moreover, the $L(u,j)$ represent all the isomorphism classes of simple $\mathcal{D}(\Lambda_{n,d})$-modules when u and j vary in $\mathbb{Z}_n$. Those of dimension d are also projective, and there are $\frac{n^2}{d}$ projective simples.

The simple modules were characterised in [Rd2], and when $n = d$, the simple modules have been described in [Ch].

We can now decompose each Γ_u, and therefore $\mathcal{D}(\Lambda_{n,d})$, into blocks:

Theorem 2.25 *The algebras Γ_u decompose into blocks as follows: if $j_{u,1}, \ldots, j_{u,r_u}$ are the representatives of the orbits of σ_u in $\mathbb{Z}_n$, then $\Gamma_u = \bigoplus_{i=1}^{r_u} \mathbb{B}_{u,i}$ where*

$$\mathbb{B}_{u,i} = \bigoplus_{t} \bigoplus_{h=0}^{d-\langle 2j+u-1\rangle^{-}-1} P(u, \sigma^t(j_{u,i}))X^h,$$

where t ranges from 0 to $\frac{2n}{d} - 1$ if $2j_{u,i} + u - 1 \not\equiv 0$ and $t = 0$ if $2j_{u,i} + u - 1 \equiv 0$.

It then follows that the quiver of $\mathcal{D}(\Lambda_{n,d})$ has $\frac{n^2}{d}$ isolated vertices which correspond to the simple projective modules, and $\frac{n(d-1)}{2}$ copies of the quiver



with $\frac{2n}{d}$ vertices and $\frac{4n}{d}$ arrows. The relations on this quiver are bb, $\overline{b}\overline{b}$ and $b\overline{b}-\overline{b}b$ (there are $\frac{6n}{d}$ relations on each of these quivers). The vertices in this quiver correspond to the simple modules $L(u,j)$, $L(u,\sigma_u(j))$, $L(u,\sigma_u^2(j))$, $\ldots$, $L(u,\sigma_u^{\frac{2n}{d}-1}(j))$.

Proof: For the general principle of presenting a basic algebra by quiver and relations, see [ARS, Section II.5 and III.1 Theorem 1.9]. The arrows of the quiver correspond to generators of $\mathrm{rad}(\mathcal{D}(\Lambda_{n,d}))/\mathrm{rad}^2(\mathcal{D}(\Lambda_{n,d}))$. From the structure of the indecomposable projective modules we know that for each p in $\mathbb{Z}_{\frac{2n}{d}}$, there is one arrow which we call b_p from the vertex ϵ_p – corresponding to the simple module $L(u,\sigma_u^p(i))$ – to the vertex ϵ_{p+1}, and one arrow, called $\overline{b}_p$, from ϵ_{p+1} to ϵ_p. The zero relations follow easily, and moreover, for each p, there is a non-zero scalar c_p with $c_p\overline{b}_pb_p = b_{p-1}\overline{b}_{p-1}$.

Starting at $p=1$, we replace $c_pc_{p-1}\ldots c_1\overline{b}_p$ by $\overline{b}_p$, for $p=1,2,\cdots,\frac{2n}{d}-1$, so that the relations become

$$\begin{array}{ll} \overline{b}_pb_p = b_{p-1}\overline{b}_{p-1} & \text{for } 1\leqslant p\leqslant \frac{2n}{d}-1 \\ c_0\ldots c_{\frac{2n}{d}-1}\overline{b}_0b_0 = b_{-1}\overline{b}_{-1}. & \end{array}$$

Since the algebra is symmetric (see the Introduction), the scalar in the final relation is 1: to see this, take a symmetrising form $\psi:\mathbb{B}_{u,i}\to k$; then

$$\psi(c_0\ldots c_{\frac{2n}{d}-1}\overline{b}_0b_0) = \psi(b_{-1}\overline{b}_{-1}) = \psi(\overline{b}_{-1}b_{-1}) = \psi(b_0\overline{b}_0) = \psi(\overline{b}_0b_0).$$

Hence $(c_0\ldots c_{\frac{2n}{d}-1}-1)\overline{b}_0b_0$ lies in the kernel of ψ, which spans a 1-dimensional (left) ideal of the algebra, and since ψ is non-singular, this must be zero. This shows that $c_0\ldots c_{\frac{2n}{d}-1}=1$.

■

Remark **2.26** It is now easy to see from the quiver and the structure of the projectives that the algebra $\mathcal{D}(\Lambda_{n,d})$ is special biserial (see [E, II.1]).

Note also that it follows from [E, II.3.1] that $\mathcal{D}(\Lambda_{n,d})$ is therefore tame or of finite type (see [ARS, p111] and [B, 4.4] for the definitions). We shall see in Section 3 and in the Appendix that it is in fact tame.

Remark **2.27** It is known that the finite-dimensional quotients of $U_q(\mathfrak{sl}_2)$ studied by R. Suter [S], J. Xiao [X] and M. Patra [P] are quotients of some of these Drinfel'd doubles (for particular choices of d). Therefore their results can be recovered from the study of the Drinfel'd doubles $\mathcal{D}(\Lambda_{n,d})$.

Remark **2.28** If we fix a block $\mathbb{B}$ of $\mathcal{D}(\Lambda_{n,d})$ which is not simple, we can prove that it is a Koszul algebra, using the results in [GM, Section 3] (by looking at minimal projective resolutions of the simple modules for $\mathbb{B}$). Therefore, by [GM, Theorem 6.1 and Section 10], its Koszul dual $\mathrm{Ext}^*_{\mathbb{B}}(\mathbb{B}/\mathrm{rad}(\mathbb{B}),\mathbb{B}/\mathrm{rad}(\mathbb{B}))$ is given by $k\mathcal{Q}^{op}/I^{\perp}$, where $\mathcal{Q}$ is the quiver of $\mathbb{B}$ and $I^{\perp}$ is generated by the relations $b\overline{b}+\overline{b}b$ for all the arrows $b,\overline{b}$ in the quiver.

In fact, we see that $\mathrm{Ext}^*_{\mathbb{B}}(\mathbb{B}/\mathrm{rad}(\mathbb{B}),\mathbb{B}/\mathrm{rad}(\mathbb{B}))$ is the preprojective algebra associated to an (unoriented) cycle. In [G, Theorem 7.2], it was shown that such an algebra is Koszul, giving another proof that $\mathbb{B}$ is a Koszul algebra.

3 Classification of the representations of $\mathcal{D}(\Lambda_{n,d})$

We can now classify all the indecomposable representations of $\mathcal{D}(\Lambda_{n,d})$ and describe its Auslander-Reiten quiver. For definitions of **Auslander-Reiten sequences** (also called **almost split sequences**) and **Auslander-Reiten quivers**, as well as descriptions of some components of Auslander-Reiten quivers, see [ARS, V.1 and VII.1] and [E, I.7 and I.8]. In this section we describe the representations and the Auslander-Reiten quiver without proofs; these are outlined in the Appendix. Note that this description shows that the algebra $\mathcal{D}(\Lambda_{n,d})$ is tame (see Remark 2.26). We then determine the quantum dimension of these representations and study related properties which we use in Section 4 when we calculate the tensor products of representations.

3.1 Description of the indecomposable modules

(I) String modules of odd length: the indecomposable modules of odd length are syzygies of simple modules, that is, of the form $\Omega^k(L(u,i))$ for some k in $\mathbb{Z}$ and for some simple module $L(u,i)$. We have

$$|\mathrm{length}(\mathrm{top}(\Omega^k(L(u,i)))) - \mathrm{length}(\mathrm{soc}(\Omega^k(L(u,i))))| = 1.$$

These modules are not periodic.

(II) String modules of even length: fix a block $\mathbb{B}_{u,i}$. For each $0 \leqslant p \leqslant \frac{2n}{d} - 1$ and for each $\ell \geqslant 1$, there are two indecomposable modules of length 2ℓ which we call $M_{2\ell}^{\pm}(u, \sigma_u^p(i))$:

- The module $M_{2\ell}^{+}(u, \sigma_u^p(i))$ has top composition factors $L(u, \sigma_u^p(i))$, $L(u, \sigma_u^{p+2}(i))$, $\dots$, $L(u, \sigma_u^{p+2(\ell-1)}(i))$ and socle composition factors $L(u, \sigma_u^{p+1}(i))$, $L(u, \sigma_u^{p+3}(i))$, $\dots$, $L(u, \sigma_u^{p+2(\ell-1)+1}(i))$:

$$\begin{tikzcd}
L(u,\sigma_u^p(i)) \arrow[dr, dash] & & L(u,\sigma_u^{p+2}(i)) \arrow[dr, dash] & & L(u,\sigma_u^{p+2(\ell-1)}(i)) \arrow[dr, dash] & \\
& L(u,\sigma_u^{p+1}(i)) \arrow[ur, dash] & & \cdots \arrow[ur, dash] & & L(u,\sigma_u^{p+2(\ell-1)+1}(i))
\end{tikzcd}$$

  The lines joining the simple modules are given by multiplication by the appropriate b-arrow or $\bar{b}$-arrow (in the case $n = d$, when there is an ambiguity, the first line is multiplication by $\gamma^{\dim L(u,\sigma_u^p(i))}$, the next one is multiplication by a scalar multiple of $X^{d-\dim L(u,\sigma_u^p(i))}$, and so on, up to scalars).

- The module $M_{2\ell}^{-}(u, \sigma_u^p(i))$ has top composition factors $L(u, \sigma_u^p(i))$, $L(u, \sigma_u^{p-2}(i))$, $\dots$, $L(u, \sigma_u^{p-2(\ell-1)}(i))$ and socle composition factors $L(u, \sigma_u^{p-1}(i))$, $L(u, \sigma_u^{p-3}(i))$, $\dots$, $L(u, \sigma_u^{p-2(\ell-1)+1}(i))$:

$$\begin{tikzcd}
& L(u,\sigma_u^{p-2(\ell-1)}(i)) \arrow[dr, dash] & & L(u,\sigma_u^{p-2}(i)) \arrow[dr, dash] & & L(u,\sigma_u^{p}(i)) \\
L(u,\sigma_u^{p-2(\ell-1)+1}(i)) \arrow[ur, dash] & & \cdots \arrow[ur, dash] & & L(u,\sigma_u^{p-1}(i)) \arrow[ur, dash] &
\end{tikzcd}$$

  As for the other string modules, the lines represent multiplication by an appropriate b or $\bar{b}$ arrow, and in the case $n = d$ the first one from the left is multiplication by a power of X and so on.

In both cases, indices are taken modulo $\frac{2n}{d}$.

These modules are periodic of period $\frac{2n}{d}$.

(III) Band modules (even length): fix a block $\mathbb{B}_{u,i}$. For each $\lambda \neq 0$ in k, and for each $\ell \geqslant 1$, there are two indecomposable modules of length $\frac{2n}{d}\ell$, which we denote by $C_\lambda^{\ell\pm}(u,i)$. They are defined as follows:

- Let V be an ℓ-dimensional vector space. Then $C_\lambda^{\ell+}(u,i)$ has underlying space $C^+ = \bigoplus_{p=0}^{\frac{2n}{d}-1} C_p^+$ with $C_p^+ = V$ for all p. The action of the idempotents ϵ_p is such that $\epsilon_p C^+ = C_p^+$. The action of the arrows $\bar{b}_{2p}$ and b_{2p+1} is zero. The action of the arrows $\bar{b}_{2p+1}$ is the identity of V. The action of the arrows b_{2p} with $p \neq 0$ is also the identity. Finally, the action of b_0 is given by the indecomposable Jordan matrix $J_\ell(\lambda)$.
  Note that $\mathrm{soc}(C^+) = \mathrm{rad}(C^+) = \bigoplus_p \epsilon_{2p+1} C^+$ and that $C^+/\mathrm{rad}(C^+) = \bigoplus_p \epsilon_{2p} C^+$.

- The module $C_\lambda^{\ell-}(u,i)$ is defined similarly, interchanging b's and $\bar{b}$'s. Note that if $C^- = \bigoplus_{p=0}^{\frac{2n}{d}-1} C_p^-$ is the underlying vector space of $C_\lambda^{\ell-}(u,i)$, the Jordan matrix $J_\ell(\lambda)$ occurs as $\bar{b}_0$ from C_1^- to C_0^-, and that we have $\mathrm{soc}(C^-) = \bigoplus_p \epsilon_{2p}C^-$ and $C^-/\mathrm{rad}(C^-) = \bigoplus_p \epsilon_{2p+1}C^-$.

These modules are periodic of period 2.

3.2 Description of the Auslander-Reiten components

(I) Components with indecomposable modules of odd length: there are two components in the Auslander-Reiten quiver for each block $\mathbb{B}_{u,i}$, one of which contains the simple modules $L(u,\sigma_u^{2p}(i))$ and the other containing the simple modules $L(u,\sigma_u^{2p+1}(i))$ for all $0 \leqslant p \leqslant \frac{2n}{d}-1$. They are of tree class $\tilde{A}_{\frac{2n}{d}-1}$, infinite in all directions, but we identify along horizontal lines so that the component lies on an infinite cylinder:



where S_p represents $L(u,\sigma_u^p(i))$, and P_p is the projective $P(u,\sigma_u^p(i))$.

(II) Components with string modules of even length: there are four components in the Auslander-Reiten quiver for each block $\mathbb{B}_{u,i}$, one which contains the modules $M_{2\ell}^+(u,\sigma_u^{2p}(i))$ for all p and ℓ, one which contains all the modules $M_{2\ell}^+(u,\sigma_u^{2p+1}(i))$ for all p and ℓ, one which contains the modules $M_{2\ell}^-(u,\sigma_u^{2p}(i))$ for all p and ℓ, and the final one contains all the modules $M_{2\ell}^-(u,\sigma_u^{2p+1}(i))$ for all p and ℓ. They are tubes of rank $\frac{n}{d}$, and the modules of length 2ℓ form the ℓ^{th} row.

The Auslander-Reiten sequences are:

$$0 \to M_{2\ell}^+(u,i-d) \to \begin{matrix} M_{2\ell+2}^+(u,i-d) \\ \oplus \\ M_{2\ell-2}^+(u,i) \end{matrix} \to M_{2\ell}^+(u,i) \to 0$$

and

$$0 \to M_{2\ell}^-(u,i+d) \to \begin{matrix} M_{2\ell+2}^-(u,i+d) \\ \oplus \\ M_{2\ell-2}^-(u,i) \end{matrix} \to M_{2\ell}^-(u,i) \to 0$$

(where $M_0^\pm(u,i) := 0$).

(III) Components with band modules: fix a block $\mathbb{B}_{u,i}$. There are two components in the Auslander-Reiten quiver for each nonzero parameter λ in k, one which contains the modules $C_\lambda^{\ell+}(u,i)$ for all ℓ and the other which contains the modules $C_\lambda^{\ell-}(u,i)$ for all ℓ. They are tubes of rank one.

The Auslander-Reiten sequences are:

$$0 \to C_\lambda^{\ell+}(u,i) \to \begin{matrix} C_\lambda^{\ell+1,+}(u,i) \\ \oplus \\ C_\lambda^{\ell-1,+}(u,i) \end{matrix} \to C_\lambda^{\ell+}(u,i) \to 0$$

and

$$0 \to C_\lambda^{\ell -}(u,i) \to \begin{array}{c} C_\lambda^{\ell+1,-}(u,i) \\ \oplus \\ C_\lambda^{\ell-1,-}(u,i) \end{array} \to C_\lambda^{\ell -}(u,i) \to 0$$

where $C_\lambda^{0\pm}(u,i) := 0$.

3.3 Splitting trace modules

In this section, we determine the quantum dimensions of the modules described above. This concept is defined in [CP] for ribbon algebras (note that the Drinfel'd double of a Hopf algebra naturally gives rise to a ribbon algebra and that $S(x) = GxG^{-1}$ for all $x \in \mathcal{D}(\Lambda_{n,d})$):

Definition 3.1 *[CP, Section 4.2.C] Let* $\rho : \mathcal{D}(\Lambda_{n,d}) \to \mathrm{End}_k(V)$ *be a representation of* $\mathcal{D}(\Lambda_{n,d})$*. Let* $f : V \to V$ *be a linear map and let* tr *be the usual trace.*

- *The quantum trace of* f *is* $\mathrm{qtr}(f) = \mathrm{tr}(\rho(G)f)$*.*
- *The quantum dimension of* V *is* $\mathrm{qdim}(V) = \mathrm{qtr}(\mathrm{id}_V)$*.*

In Section 4, we shall consider tensor products of modules over k, and in this context, **splitting trace modules,** that is, modules M such that the trivial module $L(0,0)$ is a direct summand in $\mathrm{End}_{\mathcal{D}(\Lambda_{n,d})}(M)$, are useful in view of the following proposition:

Proposition 3.2 *Let* M *be an indecomposable module, and let* $\mathcal{A}(M)$ *be its Auslander-Reiten sequence.*

(i) Suppose N *is a splitting trace module, and* $M \otimes N = \bigoplus_i A_i \oplus \bigoplus_j B_j \oplus P$*, with the* A_i *non-projective indecomposable splitting trace modules, the* B_j *non-projective indecomposable and not splitting trace modules, and* P *projective. Then* $\mathcal{A}(M) \otimes N$ *is the direct sum of* $\bigoplus_i \mathcal{A}(A_i)$ *and of a split exact sequence (equal to* $0 \to \bigoplus_j \Omega^2(B_j) \oplus Q \to \bigoplus_j \Omega^2(B_j) \oplus \bigoplus_j B_j \oplus Q \oplus P \to \bigoplus_j B_j \oplus P \to 0$ *for some projective* Q*).*

(ii) If N *is not a splitting trace module, then* $\mathcal{A}(M) \otimes N$ *is split exact.*

This proposition is proved using the same methods as [H, 2.5.9], [AC, 2.6], [GMS, 3.1 and 3.8] and [Ka, 1.5], replacing the natural isomorphism $M \cong M^{**}$ by the isomorphism of $\mathcal{D}(\Lambda_{n,d})$-modules given by $m \mapsto \langle -, Gm \rangle$.

Splitting trace modules are related to the quantum trace as follows:

Proposition 3.3 *[H, 2.5.9] A module* M *is a splitting trace module if and only if there exists an endomorphism* $f \in \mathrm{End}_{\mathcal{D}(\Lambda_{n,d})}(M)$ *such that* $\mathrm{qtr}(f) \neq 0$*.*

Remark **3.4** Note that if $\mathrm{qdim}(M) \neq 0$, then M is a splitting trace module.

Proposition 3.5 *Let* $M = \Omega^\ell(L(u,i))$ *be a module of odd length. Then* $\mathrm{qdim}(M) = (-1)^\ell q^{i+u} \dfrac{1 - q^{\dim(L(u,i))}}{1-q}$*. Therefore if* M *is not projective, then it is a splitting trace module.*

Proof: We prove it for ℓ non-negative and even; the other cases are similar:

$$\begin{tikzcd}
M = & L(u,\sigma_u^{-\ell}(i)) \arrow[dr, dash] & & L(u,\sigma_u^{\ell-2}(i)) \arrow[dr, dash] & & L(u,i) \arrow[dr, dash] & & L(u,\sigma_u^{\ell}(i)) \ . \\
& & L(u,\sigma_u^{-\ell+1}(i)) \arrow[ur, dash] & & \cdots \arrow[ur, dash] & & \cdots \arrow[ur, dash] &
\end{tikzcd}$$

Set $N = \dim L(u,i)$. For each composition factor, choose a basis obtained by taking a generator of this simple which is in the kernel of the action of X, and then applying arrows to this element. These bases can be chosen so that they are compatible with the actions of the b- and $\bar{b}$-arrows. This gives a basis for M, with respect to which the action of G can be described by the diagonal matrix with entries q^{i+u}, q^{i+u+1}, …, $q^{i+u+N-1}$, $q^{\sigma_u(i)+u}$, …, $q^{\sigma_u(i)+u+d-N-1}$, q^{i+u}, q^{i+u+1}, …, $q^{i+u+N-1}$, …, $q^{i+u+N-1}$, so that $\mathrm{qdim}(M)$, which is the trace of this matrix, is

$$\ell\left(\sum_{t=0}^{N-1} q^{i+u+t} + \sum_{s=0}^{d-N-1} q^{\sigma_u(i)+u+s}\right) + \sum_{t=0}^{N-1} q^{i+u+t}.$$

Since $\sigma_u(i) = i+N$, setting $s = t-N$ gives:

$$\mathrm{qdim}(M) = \ell\sum_{t=0}^{d-1} q^{i+u+t} + \sum_{t=0}^{N-1} q^{i+u+t} = \ell q^{i+u}\frac{1-q^d}{1-q} + q^{i+u}\frac{1-q^N}{1-q} = q^{i+u}\frac{1-q^N}{1-q}.$$

Therefore M is a splitting trace module if, and only if, $N \neq d$, that is, M is not projective.

■

Proposition 3.6 *Let M be an indecomposable module of even length. Then M is not a splitting trace module and* $\mathrm{qdim}(M) = 0$.

Proof: Take f in $\mathrm{End}_{\mathcal{D}(\Lambda_{n,d})}(M)$. We choose a basis for M as in the proof of Proposition 3.5. Since f is a homomorphism of $\mathcal{D}(\Lambda_{n,d})$-modules, the matrix of f with respect to this basis is diagonal. By considering the actions of the arrows b_p and $\bar{b}_p$ which connect the composition factors of M, we can see that f is of the form $\mu\mathrm{id}$ for some $\mu \in k$. Therefore $\mathrm{qtr}(f) = \mu\mathrm{qtr}(\mathrm{id}) = \mu\mathrm{qdim}(M)$. We can then calculate this as in the proof of Proposition 3.5, to see that $\mathrm{qdim}(M) = 0$. Therefore $\mathrm{qtr}(f) = 0$ for all $f \in \mathrm{End}_{\mathcal{D}(\Lambda_{n,d})}(M)$ and so M is not a splitting trace module. ■

4 Tensor products of $\mathcal{D}(\Lambda_{n,d})$-modules

In this section, we determine the tensor products over the base field k of modules of types **(I)** and **(II)**. Recall that if M and N are modules over $\mathcal{D}(\Lambda_{n,d})$, then $M \otimes N \cong N \otimes M$ (where $\otimes = \otimes_k$).

4.1 Tensor product of simple $\mathcal{D}(\Lambda_{n,d})$-modules

The aim of this section is to prove the following theorem:

Theorem 4.1 *Set $\varpi = \dim L(u,i) + \dim L(v,j) - (d+1)$, and define*

$$\varsigma = \begin{cases} \frac{\varpi}{2} & \text{if } \varpi \text{ is even} \\ \frac{\varpi+1}{2} & \text{if } \varpi \text{ is odd.} \end{cases}$$

Then we have the following decompositions:

(a) If $\varpi \leqslant 0$,

$$L(u,i) \otimes L(v,j) \cong \bigoplus_{\theta=0}^{\min\{\dim L(u,i)-1;\dim L(v,j)-1\}} L(u+v, i+j+\theta).$$

(b) If $\varpi \geqslant 0$,

$$L(u,i) \otimes L(v,j) \cong \bigoplus_{\theta=\varpi+1}^{\min\{\dim L(u,i)-1;\dim L(v,j)-1\}} L(u+v, i+j+\theta) \oplus \bigoplus_{\theta=\varsigma}^{\varpi} P(u+v, i+j+\theta).$$

Remark **4.2** In the case $n = d$, H-X. Chen gives this decomposition when $\varpi \leqslant 0$, and gives the socle of this tensor product when $\varpi \geqslant 0$ in [Ch].

4.1.1 Preliminaries

Recall (Proposition 2.20) that a simple module is determined up to isomorphism by its dimension, the action of the idempotents E_u and the action of the idempotents e_i. This will be important in order to determine the socle of the tensor product $L(u,i) \otimes L(v,j)$.

Notation **4.3** To simplify notation, set

$$\begin{cases} \dim L(u,i) = \alpha, \\ \dim L(v,j) = \beta, \\ \text{assume that } \alpha \leqslant \beta \text{ throughout.} \end{cases}$$

Lemma 4.4 *The vector space* $L(u,i) \otimes L(v,j)$ *has basis* $\{z_{s,t} \mid 0 \leqslant s \leqslant \alpha - 1,\ 0 \leqslant t \leqslant \beta - 1\}$, *where* $z_{s,t} = \gamma_i^s \tilde{H}_{u,i} \otimes \gamma_j^t \tilde{H}_{v,j}$.

Definition 4.5 *We define a* **grading** *on the vector space* $L(u,i) \otimes L(v,j)$ *by setting* $\deg_{\otimes}(z_{s,t}) = s + t$.

Lemma 4.6 *The socle of* $L(u,i) \otimes L(v,j)$ *is isomorphic to a direct sum of simple modules of the form* $L(u+v,-)$.

Proof: It is easy to see that $E_{u+v} z_{s,t} = z_{s,t}$ for all s,t. ■

In order to find all the simple modules in the socle, we need to find the kernel of the action of X: the generators $\tilde{H}_{w,\ell}$ of the simple modules are in the kernel of the action of X. Note that $X \cdot \gamma_\ell^s \tilde{H}_{w,\ell} = (s)_q (1 - q^{2\ell + w - 1 + t}) \gamma_\ell^{s-1} \tilde{H}_{w,\ell}$, which is nonzero if $s > 0$, and zero if $s = 0$.

4.1.2 Kernel of X

Lemma 4.7 *The kernel of the action of* X *on* $\mathcal{D}(\Lambda_{n,d})$ *has dimension* $n^2 d$.

Proof: The Hopf algebra $\Lambda_{n,d}^{*cop}$ is a Hopf subalgebra of $\mathcal{D}(\Lambda_{n,d})$, so by a theorem of Nichols-Zoeller [NZ, Theorem 7], $\mathcal{D}(\Lambda_{n,d})$ is free as a module over $\Lambda_{n,d}^{*cop}$, with rank nd.

Now $\Lambda_{n,d}^{*cop} = \langle G, X \mid G^n = 1, X^d = 0, GX = q^{-1} XG \rangle$ so the kernel of the action of X on $\Lambda_{n,d}^{*cop}$ has dimension n with basis $\{G^i X^{d-1} \mid i = 0, \ldots, n-1\}$.

Therefore the kernel of the action of X on $\mathcal{D}(\Lambda_{n,d})$ has dimension $n \cdot nd$. ■

Lemma 4.8 *The kernel of* X *is spanned by* $\{\tilde{H}_{u,j} X^h, \tilde{D}_{u,j} X^h \mid u,j \in \mathbb{Z}_n, 0 \leqslant h \leqslant d - \langle 2j + u - 1 \rangle^-\}$ *(this set contains repetitions).*

Proof: Note that all the elements $\tilde{H}_{u,j} X^h$ and $\tilde{D}_{u,j} X^h$ are in the kernel of the action of X.

We consider the case where d is odd; the case d even is similar, but the simple projective modules are distributed differently.

There are n simple projective modules of dimension d in Γ_u (there are $\frac{n}{d}$ isomorphism classes of simple projective modules, and the number of isomorphic copies of each one is d) and each one gives one element of the form $\tilde{H}_{u,j} X^h = \tilde{D}_{u,j} X^h$. The other projective modules in Γ_u have dimension $2d$, so there are $\frac{\dim \Gamma_u - n \cdot d}{2d} = \frac{n(d-1)}{2}$ of them, and each one gives two elements in the kernel of X. Therefore we have $n \cdot (n + 2\frac{n(d-1)}{2}) = n^2 d$ distinct elements in $\{\tilde{H}_{u,j} X^h, \tilde{D}_{u,j} X^h \mid u,j \in \mathbb{Z}_n, 0 \leqslant h \leqslant d - \langle 2j + u - 1 \rangle^-\}$ which are in the kernel of X and linearly independent. We can visualise these elements in the basis of the action of X as in the picture below:





Note that there are n^2d+n^2 elements in $\{\tilde{H}_{u,j}X^h, \tilde{D}_{u,j}X^h \mid u, j \in \mathbb{Z}_n, 0 \leqslant h \leqslant d - \langle 2j+u-1\rangle^-\}$, which is consistent with the fact that the elements in the kernel of the action of X coming from simple projective modules appear twice. ■

From now on, we consider modules up to isomorphism, so assume that $h = 0$.

Proposition 4.9 *The kernel of the action of X on $L(u,i)\otimes L(v,j)$ has dimension $\alpha = \dim L(u,i)$, and is spanned by elements x_θ with $\deg_\otimes x_\theta = \theta$ and $0 \leqslant \theta \leqslant \alpha - 1$ (there is one element (up to scalars) in the kernel for each degree).*

Proof: Consider the vector space U_θ spanned by the elements $z_{s,t}$ with $\deg_\otimes(z_{s,t}) = \theta$. The kernel of the action of X on $L(u,i)\otimes L(v,j)$ is the span of the kernels of the φ_θ where $\varphi_\theta : U_\theta \to U_{\theta-1}$ is right multiplication by X. It is easy to see that $\varphi_\theta(z_{s,t}) = \mu_{s,t}z_{s-1,t} + \nu_{s,t}z_{s,t-1}$ with $\mu_{s,t} = 0$ if and only if $s = 0$ and $\nu_{s,t} = 0$ if and only if $t = 0$.

First assume that $0 \leqslant \theta \leqslant \alpha - 1$: then $\dim U_\theta = \theta + 1$ and $\dim U_{\theta-1} = \theta$. The matrix of φ_θ is of the form

$$\left(\begin{array}{c|cccc} \mu & \nu & & & \\ & \mu & \ddots & 0 & \\ & 0 & \ddots & \nu & \\ & & & \mu & \nu \end{array}\right)$$

so φ_θ is onto; therefore its kernel has dimension 1 and is spanned by an element x_θ with $\deg_\otimes(x_\theta) = \theta$.

If $\theta \geqslant \alpha$, then $\dim U_{\theta-1} \geqslant \dim U_\theta$ and the matrix of φ_θ is either square (if $\alpha \leqslant \theta \leqslant \beta - 1$) or rectangular with one more line than it has columns. It contains a maximal square matrix of the form

$$\begin{pmatrix} \nu & & & \\ \mu & \ddots & 0 & \\ 0 & \ddots & \nu & \\ & & \mu & \nu \end{pmatrix}$$

so the rank of φ_θ is the dimension of U_θ, and φ_θ is therefore injective. ■

Remark **4.10** An element x_θ is either of the form $\tilde{H}_{u+v,p}$, in which case it generates a simple module, or of the form $\tilde{D}_{u+v,p}$, in which case it generates a non-simple module which cannot be a summand in the socle of $L(u,i)\otimes L(v,j)$.

Remark **4.11** Since $e_{i+j+\theta}z_{s,\theta-s} = z_{s,\theta-s}$ for all s, we also have $e_{i+j+\theta}x_\theta = x_\theta$. In particular, if x_θ generates a simple module, then this simple module must be $L(u+v, i+j+\theta)$.

4.1.3 Proof of the decomposition when $\alpha + \beta \leqslant d + 1$, and some general results

Proposition 4.12 *Suppose $\varpi < 0$ and $0 \leqslant \theta < \alpha$ or $\varpi \geqslant 0$ and $0 \leqslant \varpi < \theta < \alpha$. Then x_θ generates a simple module of dimension $< d$. The element x_θ for $2\theta = \varpi$ generates a simple projective module if and only if $\varpi > 0$ is even or $\varpi = 0$.*

Therefore $L(u,i)\otimes L(v,j)$ contains $\bigoplus_{\theta=0}^{\alpha-1} L(u+v, i+j+\theta)$ if $\varpi < 0$ and contains $\bigoplus_{\theta=\varpi+1}^{\alpha-1} L(u+v, i+j+\theta) \oplus L(u+v, i+j+\frac{\varpi}{2})$ if $\varpi \geqslant 0$, the last term only occurring if ϖ is even.

Proof: If x_θ is in the kernel of $\gamma_{i+j+\theta}^{d-1}$, then it must generate a simple module of dimension $< d$ (see Proposition 2.16). So suppose for a contradiction that $\gamma_{i+j+\theta}^{d-1}x_\theta \neq 0$. Then $\gamma_{i+j+\theta}^{d-1}z_{s,\theta-s} \neq 0$ for some s. But $\gamma_{i+j+\theta}^{d-1}z_{s,\theta-s} = \sum_{r=0}^{d-1}\binom{d-1}{r}_q q^{r(\theta-s)}\gamma_i^{r+s}\tilde{H}_{u,i}\otimes\gamma_j^{\theta+d-1-r-s}\tilde{H}_{v,j}$, and the coefficients are all nonzero. Therefore there exists an r such that $\gamma_i^{r+s}\tilde{H}_{u,i}\otimes\gamma_j^{\theta+d-1-r-s}\tilde{H}_{v,j} \neq 0$. In particular, $\gamma_i^{r+s}\tilde{H}_{u,i}$ and $\gamma_j^{\theta+d-1-r-s}\tilde{H}_{v,j}$ are nonzero, so $r+s \leqslant \alpha - 1$ and $\theta + d - 1 - r - s \leqslant \beta - 1$. Therefore $(r+s) + (\theta + d - 1 - r - s) \leqslant \alpha + \beta - 2$.

On the other hand, $(r+s)+(\theta+d-1-r-s)=\theta+d-1$.

Now consider the case $\varpi<\theta<\alpha$: then

$$\begin{aligned}\alpha+\beta-2 &= \varpi+d+1-2\\ &< \theta+d-1\\ &= (r+s)+(\theta+d-1-r-s)\\ &\leqslant \alpha+\beta-2,\end{aligned}$$

and this gives a contradiction, so x_θ must generate a simple module of dimension $<d$.

Finally, assume that $\theta=\frac{\varpi}{2}$ with ϖ even. Then either $\varpi>0$ and $2\theta=\varpi=\alpha+\beta-d-1$, or $\varpi=0$ and $\theta=0$ (since $\varpi\leqslant 0$ and $\theta=\frac{\varpi}{2}\geqslant 0$).

Suppose for a contradiction that x_θ generates a non-simple module. Then the degree of x_θ must be (see Proposition 2.16)

$$\begin{aligned}-\langle 2(i+j)+(u+v)-1+2\theta\rangle &= -\langle(2i+u-1)+(2j+v-1)+1+\varpi\rangle\\ &= -\langle d-\alpha+d-\beta+1+\varpi\rangle\\ &= -\langle d\rangle=-d,\end{aligned}$$

which is impossible since the smallest possible degree is $-d+1$.

Therefore x_θ also generates a simple module, and this module is $L(u+v,i+j+\frac{\varpi}{2})$, of dimension d and so projective.

The sum of all the simple modules above is contained in $L(u,i)\otimes L(v,j)$, and since the simple modules which occur are non-isomorphic, the sum is direct. ■

We can now give the decomposition of $L(u,i)\otimes L(v,j)$ when $\dim L(u,i)+\dim L(v,j)\leqslant d+1$:

Proposition 4.13 *Assume that* $\dim L(u,i)+\dim L(v,j)\leqslant d+1$. *Then*

$$L(u,i)\otimes L(v,j)\cong\bigoplus_{\theta=0}^{\min\{\dim L(u,i)-1,\dim L(v,j)-1\}}L(u+v,i+j+\theta).$$

Proof: We know that $L(u,i)\otimes L(v,j)$ contains $\bigoplus_{\theta=0}^{\alpha-1}L(u+v,i+j+\theta)$. Computing the dimension of this direct sum gives:

$$\dim\left(\bigoplus_{\theta=0}^{\alpha-1}L(u+v,i+j+\theta)\right)=\sum_{\theta=0}^{\alpha-1}\left(d-\langle-\alpha-\beta+1+2\theta\rangle^-\right).$$

But we have

$$-\alpha-\beta+1+2\theta\geqslant-(d+1)+1+2\theta\geqslant-d$$

and

$$-\alpha-\beta+1+2\theta\leqslant-\alpha-\beta+1+2(\alpha-1)=\alpha-\beta-1<0.$$

Therefore

$$\begin{aligned}\dim\left(\bigoplus_{\theta=0}^{\alpha-1}L(u+v,i+j+\theta)\right) &= \sum_{\theta=0}^{\alpha-1}(d-(-\alpha-\beta+1+2\theta+d))\\ &= (\alpha+\beta-1)\alpha-2\tfrac{(\alpha-1)\alpha}{2}\\ &= \alpha\beta\\ &= \dim L(u,i)\otimes L(v,j)\end{aligned}$$

■

4.1.4 Socle of $L(u,i)\otimes L(v,j)$

Proposition 4.14 *The socle of* $L(u,i)\otimes L(v,j)$ *is isomorphic to*

$$\bigoplus_{\theta=\varsigma}^{\min\{\dim L(u,i)-1;\dim L(v,j)-1\}}L(u+v,i+j+\theta).$$

The proof follows from the next two lemmas; we first consider the case $0 \leqslant 2\theta < \varpi$:

Lemma 4.15 *Assume that $0 \leqslant 2\theta < \varpi$. Then x_θ does not generate a simple module.*

Proof: Note that we have $\varpi > 0$ and therefore $\alpha + \beta > d + 1$. Suppose for a contradiction that x_θ does generate a simple module. Then it would have dimension $d - \langle -\alpha - \beta + 1 + 2\theta \rangle^- = d - (-\alpha - \beta + 1 + 2\theta + 2d) = \alpha + \beta - d - 1 - 2\theta$. Therefore, the element $\gamma^{\alpha+\beta-d-1-2\theta} x_\theta$ would be in the kernel of all arrows.

But in the same way that we determined the kernel of the action of X (see Proposition 4.9), we can see that the intersection of the kernels of the actions of all the arrows on $L(u,i) \otimes L(v,j)$ has dimension α and has a basis formed of elements y_π for $\beta - 1 \leqslant \pi \leqslant \alpha + \beta - 2$, with $\deg_\otimes y_\pi = \pi$.

Therefore, we see that the element $\gamma^{\alpha+\beta-d-1-2\theta} x_\theta$ must have a degree in $L(u,i) \otimes L(v,j)$ which is at least $\beta - 1$. However, $\deg_\otimes \gamma^{\alpha+\beta-d-1-2\theta} x_\theta = \alpha + \beta - d - 1 - \theta = \beta - 1 - (d - \alpha + \theta) < \beta - 1$. Therefore x_θ cannot generate a simple module. ■

The remaining case is $\varpi < 2\theta \leqslant 2\varpi$:

Lemma 4.16 *The socle of the module generated by x_θ with $0 \leqslant 2\theta < \varpi$ is a simple module, generated by an element $x_{\theta'}$ with $\varpi < 2\theta' \leqslant 2\varpi$, and all the remaining elements in the kernel of the action of X can be obtained in this way.*

Proof: The element x_θ is of the form $\tilde{D}_{u+v,p}$ for some p. To find p, note that $\deg_\otimes x_\theta = i + j + \theta$, so $e_{i+j+\theta}$ is the unique vertex acting as the identity on $\tilde{D}_{u+v,p}$, and therefore $i + j + \theta = p - \langle 2p + u + v - 1 \rangle = \sigma_{u+v}(p) - d$. So we have $p = \sigma^2_{u+v}(p) - d = \sigma_{u+v}(d + i + j + \theta) - d$.

We can now compute the dimension of $\frac{\Gamma_u x_\theta}{\mathrm{soc}\Gamma_u x_\theta}$; this is equal to $\langle 2p + u + v - 1 \rangle = \varpi - 2\theta$ using Proposition 2.16 and the previous remarks.

Now consider $\gamma^{i+j+\theta} x_\theta$: we know that this is an element in the kernel of the action of X. Set $\theta' := \deg_\otimes \gamma^{i+j+\theta} x_\theta = \varpi - \theta$. Then $2\theta' = \varpi + (\varpi - 2\theta) > \varpi$, so $\gamma^{i+j+\theta} x_\theta = x_{\theta'}$ up to scalars, with $\varpi < 2\theta' \leqslant 2\varpi$.

The final claim follows from the fact that $\#\{\theta \mid 0 \leqslant 2\theta < \varpi\} = \#\{\theta' \mid \varpi < 2\theta' \leqslant 2\varpi\}$. ■

4.1.5 Proof of the decomposition when $\alpha + \beta > d + 1$

We now know the socle of $L(u,i) \otimes L(v,j)$, and we want to determine the entire decomposition. For this we determine a largest semisimple summand which has no projective summands and prove that the remaining summands must be projective.

Consider the projective module $P(v, \sigma_v^{2p+1}(j))$:

$$\begin{tikzcd}
 & L(v,\sigma_v^{2p+1}(j)) \arrow[dl, dash] \arrow[dr, dash] & \\
L(v,\sigma_v^{2p}(j)) \arrow[dr, dash] & & L(v,\sigma_v^{2p+2}(j)) \arrow[dl, dash] \\
 & L(v,\sigma_v^{2p+1}(j)) &
\end{tikzcd}$$

We have an exact sequence

$$0 \to L(v,\sigma_v^{2p+1}(j)) \longrightarrow \Omega(L(v,\sigma_v^{2p+1}(j))) \longrightarrow L(v,\sigma_v^{2p}(j)) \oplus L(v,\sigma_v^{2p+2}(j)) \to 0.$$

Note that $\dim L(v,\sigma_v^{2p+1}(j)) = d - \beta < \alpha$ since $\alpha + \beta > d + 1$. Moreover, $\dim L(u,i) + \dim L(v,\sigma_v^{2p+1}(j)) = \alpha + d - \beta < d + 1$ since $\alpha \leqslant \beta$. Therefore, by Proposition 4.13, taking the direct sum of these sequences over p and tensoring on the left by $L(u,i)$ gives:

$$\begin{aligned} 0 \to \bigoplus_{p=0}^{\frac{n}{d}-1} \bigoplus_{\theta=0}^{d-\beta-1} L(u+v, i + \sigma_v^{2p+1}(j) + \theta) \xrightarrow{f} & \bigoplus_{p=0}^{\frac{n}{d}-1} \bigoplus_{\theta=0}^{d-\beta-1} \Omega(L(u+v, i + \sigma_v^{2p+1}(j) + \theta)) \oplus P \\ & \longrightarrow \left(\bigoplus_{p=0}^{\frac{n}{d}-1} L(u,i) \otimes L(v,\sigma_v^{2p}(j)) \right)^2 \to 0, \end{aligned}$$

where P is a projective module.

Set $M := \bigoplus_{p=0}^{\frac{n}{d}-1} L(u,i) \otimes L(v, \sigma_v^{2p}(j))$. We know by Proposition 4.14 that the socle of M is $\bigoplus_{p=0}^{\frac{n}{d}-1} \bigoplus_{\theta=\varsigma}^{d-\alpha-1} L(u+v, i+\sigma_v^{2p}(j)+\theta)$. It is straightforward to check that the summands in $\bigoplus_{\theta=0}^{d-\beta-1} L(u+v, i+\sigma_v^{2p+1}(j)+\theta)$ and in $\mathrm{soc}(M)$ are pairwise non-isomorphic.

We now determine the nonprojective part of M.

Take a simple module $S := L(u+v, i+\sigma_v^{2p+1}(j)+\theta)$ inside the left-hand term. If S is projective, then it must be a summand in P (since $\Omega(S) = 0$, or since S is also injective), and does not occur in M.

Now assume that S is not projective. This simple module must embed via f into either

$$\begin{tikzcd}
L(u+v,\sigma_{u+v}^{-1}(i+\sigma_v^{2p+1}(j)+\theta)) & & L(u+v,\sigma_{u+v}(i+\sigma_v^{2p+1}(j)+\theta)) \\
& S \arrow[ul, dash] \arrow[ur, dash] &
\end{tikzcd}$$

or an indecomposable projective summand Q of P. Assume for a contradiction that S embeds into a projective Q. Then the cokernel of this embedding is

$$Q/S\colon \quad \begin{tikzcd}
& S \arrow[dl, dash] \arrow[dr, dash] & \\
L(u+v,\sigma_{u+v}^{-1}(i+\sigma_v^{2p+1}(j)+\theta)) & & L(u+v,\sigma_{u+v}(i+\sigma_v^{2p+1}(j)+\theta))
\end{tikzcd} .$$

This is therefore an indecomposable summand of M^2, and hence of M. So the square of Q/S must be a summand of M^2. The second copy of Q/S must arise in the same way, so there are two copies of the simple module S in $\bigoplus_{\theta=0}^{d-\beta-1} L(u+v, i+\sigma_v^{2p+1}(j)+\theta)$, which contradicts the fact that the summands are pairwise non-isomorphic.

Therefore S embeds in

$$\begin{tikzcd}
L(u+v,\sigma_{u+v}^{-1}(i+\sigma_v^{2p+1}(j)+\theta)) & & L(u+v,\sigma_{u+v}(i+\sigma_v^{2p+1}(j)+\theta)) \\
& L(u+v,i+\sigma_v^{2p+1}(j)+\theta) \arrow[ul, dash] \arrow[ur, dash] &
\end{tikzcd} .$$

So we have $M^2 \cong P \oplus \bigoplus_{p=0}^{\frac{n}{d}-1} \bigoplus_{\theta=0}^{d-\beta-1} L(u+v, \sigma_{u+v}(i+\sigma_v^{2p+1}(j)+\theta)) \oplus \bigoplus_{p=0}^{\frac{n}{d}-1} \bigoplus_{\theta=0}^{d-\beta-1} L(u+v, \sigma_{u+v}^{-1}(i+\sigma_v^{2p+1}(j)+\theta))$ which is isomorphic to $\bigoplus_{p=0}^{\frac{n}{d}-1} \bigoplus_{\theta=0}^{d-\beta-1} L(u+v, \sigma_{u+v}(i+\sigma_v^{2p+1}(j)+\theta))^2$ (recall that $\sigma^2(\ell) = \ell + d$).

We know the socle of M, so we need to identify which simple modules in the socle we have obtained. We can see that $\sigma_{u+v}(i+\sigma_v^{2p+1}(j)+\theta) = i+\sigma_v^{2p+2}(j)+(\alpha-1-\theta)$, and $\varpi + 1 = \alpha+\beta-d \leqslant \pi := \alpha-1-\theta \leqslant \alpha-1$. So $M \cong \bigoplus_{p=0}^{\frac{n}{d}-1} \bigoplus_{\pi=\varpi+1}^{\alpha-1} L(u+v, i+\sigma_v^{2p}(j)+\pi) \oplus P'$ with P' projective. Since we know the socle of M, we get $P' \cong \bigoplus_{p=0}^{\frac{n}{d}-1} \bigoplus_{\pi=\varsigma}^{\varpi} P(u+v, i+\sigma_v^{2p}(j)+\pi)$.

Finally, since all the summands in M are non-isomorphic, we conclude using the socle of $L(u,i) \otimes L(v,j)$.

4.2 Tensor product of modules of odd length

We know from the description of the modules of odd length 3.1**(I)** that all the modules of odd length are syzygies of simple modules. Moreover, we have

$$\begin{aligned}
\Omega^k(M) \otimes \Omega^\ell(N) &\cong \Omega^k(M \otimes \Omega^\ell(N)) \oplus \text{ (projectives)} \\
&\cong \Omega^k(\Omega^\ell(N) \otimes M) \oplus \text{ (projectives)} \\
&\cong \Omega^k(\Omega^\ell(N \otimes M)) \oplus \text{ (projectives)} \\
&\cong \Omega^{k+\ell}(M \otimes N) \oplus \text{ (projectives)}.
\end{aligned}$$

Therefore, from the results above applied with M and N simple, we can decompose the tensor product of any modules of odd length into a direct sum of modules of odd length, up to projectives.

4.3 Tensor product of a string module of even length with a module of odd length

In this section, we study the tensor product of a string module of even length with a module of odd length, and by 3.1**(I)** and Section 4.2 it is enough to consider the tensor product of a string module of even length with a simple module. To simplify notations, let $\mathfrak{I}$ denote the set $\{0 \leqslant \theta \leqslant \min\{\dim L(u,i), \dim L(v,j)\} - 1\}$ if $\dim L(u,i) + \dim L(v,j) < d+1$, and the set $\{\varpi + 1 \leqslant \theta \leqslant \min\{\dim L(u,i), \dim L(v,j)\} - 1\}$ if $\varpi := \dim L(u,i) + \dim L(v,j) - (d+1) \geqslant 0$.

Proposition 4.17 *The tensor product of a string module of length two with a simple module decomposes as follows:*

$$M_2^+(u,i) \otimes L(v,j) \cong \bigoplus_{\theta \in \mathfrak{I}} M_2^+(u+v, i+j+\theta) \oplus \text{projective}$$

and

$$M_2^-(u,i) \otimes L(v,j) \cong \bigoplus_{\theta \in \mathfrak{I}} M_2^-(u+v, i+j+\theta) \oplus \text{projective}.$$

Proof: We prove it for $M_2^+(u,i)$. Consider the exact sequence

$$0 \to L(u, \sigma_u(i)) \to M_2^+(u,i) \to L(u,i) \to 0$$

and tensor it with $L(v,j)$. We know the decomposition of the two outside terms from Theorem 4.1, so $M_2^+(u,i) \otimes L(v,j) = W \oplus P$ where P is a projective module and the composition factors of W are the non-projective simple summands in $L(u,i) \otimes L(v,j)$ and $L(u,\sigma_u(i)) \otimes L(v,j)$, which are known.

We now show that the non-projective summands in $M_2^+(u,i) \otimes L(v,j)$ have length 2. We know that $M_2^+(u,i)$ is periodic (from 3.1(**II**)), so $\Omega^m(M_2^+(u,i)) \cong M_2^+(u,i)$ for some $m \in \mathbb{Z}$. Then, tensoring a minimal projective resolution of $M_2^+(u,i)$ by $L(v,j)$ and using the fact that $P \otimes L(v,j)$ is projective if P is projective [GMS, Proposition 2.1] gives $\Omega^m(M_2^+(u,i) \otimes L(v,j)) \cong M_2^+(u,i) \otimes L(v,j) \oplus$ projective. So any summand of $M_2^+(u,i) \otimes L(v,j)$ is either periodic, and therefore of even length, or projective. From the description of the indecomposable modules of even length (3.1), we know that all the simple summands in the top have the same dimension N and all the simple summands in the socle have the same dimension $d - N$. It is easy to check that the (non-projective) simple modules which are summands in $L(u,i) \otimes L(v,j)$ all have different dimensions, and that for each dimension N that occurs there is exactly one simple summand in $L(u,\sigma_u(i)) \otimes L(v,j)$ which has dimension $d - N$, and moreover that the simple modules pair off to give indecomposable summands of length 2, whose top and socle are fully determined by the decompositions of $L(u,i) \otimes L(v,j)$ and $L(u,\sigma_u(i)) \otimes L(v,j)$.

We now need to decide whether these summands of length 2 are string or band modules. If $n \neq d$, then there are no band modules of length 2 so it is clear.

Now assume that $n = d$, and let us write temporarily $C_0^{1+}(u,i)$ for $M_2^+(u,i)$. For any $\lambda \in k$, the module $C_\lambda^{1+}(u,i)$ is characterised by the following pullback diagram:

$$\begin{tikzcd}
0 \arrow[r] & L(u,\sigma_u(i)) \arrow[r] \arrow[d, equal] & C_\lambda^{1+}(u,i) \arrow[r] \arrow[d] & L(u,i) \arrow[r] \arrow[d, "\varphi_\lambda"] & 0 \\
0 \arrow[r] & L(u,\sigma_u(i)) \arrow[r] & P(u,\sigma_u(i)) \arrow[r] & \Omega^{-1}(L(u,\sigma_u(i))) \arrow[r] & 0
\end{tikzcd}$$

with $\varphi_\lambda(\tilde{H}_{u,i}) = \lambda \bar{b}_0 D_{u,\sigma_u(i)} + b_1 D_{u,\sigma_u(i)}$.

The summands in $C_0^{1+}(u,i) \otimes L(v,j)$ are of the form $C_\lambda^{1+}(u+v, i+j+\theta)$. Consider such a summand: then $\varphi_\lambda : L(u+v, i+j+\theta) \to \Omega(L(u+v, \sigma_{u+v}(i+j+\theta)))$ is the restriction of $\varphi_0 \otimes id : L(u,i) \otimes L(v,j) \to \Omega(L(u,i)) \otimes L(v,j)$ to the summand $L(u+v, i+j+\theta)$. Applying b_0 to the first term in $\varphi_\lambda(\tilde{H}_{u+v,i+j+\theta})$ gives $\lambda b_0 \bar{b}_0 D_{u+v,i+j+\theta}$, and applying $\bar{b}_1$ to the second term gives $\bar{b}_1 b_1 D_{u+v,i+j+\theta} = b_0 \bar{b}_0 D_{u+v,i+j+\theta}$. So the first one is λ times the second, and they are both nonzero. Now apply the same procedure with $\varphi_0 \otimes \mathrm{id}$: the first term will be 0, and the second non-zero. Hence we must have $\lambda = 0$ and therefore the summands in $M_2^+(u,i) \otimes L(v,j)$ which are of the form $C_\lambda^{1+}(u+v, i+j+\theta)$ are in fact of the form $M_2^+(u+v, i+j+\theta)$.

We then consider all the cases:

- $\dim L(u,i) + \dim L(v,j) < d+1$ and $\dim L(u,i) \leqslant \dim L(v,j)$;

- $\dim L(u,i) + \dim L(v,j) < d+1$ and $\dim L(u,i) > \dim L(v,j)$;

- $\dim L(u,i) + \dim L(v,j) \geqslant d+1$ and $\dim L(u,i) > \dim L(v,j)$;
- $\dim L(u,i) + \dim L(v,j) \geqslant d+1$ and $\dim L(u,i) \leqslant \dim L(v,j)$

in order to determine the exact bounds (for θ) of the decomposition up to projectives.

■

Theorem 4.18 *The tensor product of a string module of even length with a simple module decomposes as follows:*

$$M^+_{2\ell}(u,i) \otimes L(v,j) \cong \bigoplus_{\theta \in \mathfrak{I}} M^+_{2\ell}(u+v, i+j+\theta) \oplus \text{projective}$$

and

$$M^-_{2\ell}(u,i) \otimes L(v,j) \cong \bigoplus_{\theta \in \mathfrak{I}} M^-_{2\ell}(u+v, i+j+\theta) \oplus \text{projective}$$

Proof: We work by induction on ℓ, using Proposition 3.2 (i), which we apply here with $M = M^+_{2\ell}(u, i+d)$ and $N = L(v,j)$. We know that N is a splitting trace module (Proposition 3.5) and we have the Auslander-Reiten sequence

$$\mathcal{A}(M^+_{2\ell}(u,i+d)): \quad 0 \to M^+_{2\ell}(u,i) \to \begin{array}{c} M^+_{2\ell+2}(u,i) \\ \oplus \\ M^+_{2\ell-2}(u,i+d) \end{array} \to M^+_{2\ell}(u,i+d) \to 0$$

(where $M^+_0(u,i) := 0$). The summands in $M^+_{2\ell+2}(u,i+d) \otimes L(v,j)$ are not splitting trace modules (they have even length), therefore, by Proposition 3.2 (i), the sequence $\mathcal{A}(M^+_{2\ell}(u,i+d)) \otimes L(v,j)$ is split exact. We know the decomposition of the tensor product of three of the terms with $L(v,j)$, so induction yields the fourth. ■

Remark **4.19** We can now find the tensor product of any string module of even length with any string module of odd length up to projectives, using the results and methods of the previous sections.

4.4 Tensor product of string modules of even length

Proposition 4.20 *The tensor product of two string modules of length two of the same type decomposes as follows:*

$$M^+_2(u,i) \otimes M^+_2(v,j) \cong \bigoplus_{\theta \in \mathfrak{I}} [M^+_2(u+v, i+j+\theta) \oplus M^+_2(u+v, \sigma_{u+v}(i+j+\theta))] \oplus \text{projective}$$

and

$$M^-_2(u,i) \otimes M^-_2(v,j) \cong \bigoplus_{\theta \in \mathfrak{I}} [M^-_2(u+v, i+j+\theta) \oplus M^-_2(u+v, \sigma^{-1}_{u+v}(i+j+\theta))] \oplus \text{projective.}$$

Proof: We start with the exact sequence $0 \to L(v, \sigma_v^{-1}(j)) \to \Omega^{-1}(L(v,j)) \to M^+_2(v,j) \to 0$. Tensoring on the left by $M^+_2(u,i)$ gives again an exact sequence, and using Proposition 4.17 we get the exact sequence:

$$0 \to \oplus_\rho M^+_2(u+v, i+\sigma_v^{-1}(j+\rho)) \to \Omega^{-1}(\oplus_\theta M^+_2(u+v,i+j+\theta)) \oplus P \to M^+_2(u,i) \oplus M^+_2(v,j) \to 0$$

where P is a projective module. Note that the middle term is equal to $\oplus_\theta M^+_2(u+v, \sigma_{u+v}(i+j+\theta)) \oplus P$ and that $i + \sigma_v^{-1}(j) + \rho = \sigma^{-1}_{u+v}(i+j+\pi)$ with π in the same range as θ (for the latter, we consider all the four cases that occurred in the proof of Proposition 4.17). So we have

$$0 \to \oplus_\pi M^+_2(u+v, \sigma^{-1}_{u+v}(i+j+\pi)) \to \oplus_\theta M^+_2(u+v, \sigma_{u+v}(i+j+\theta)) \oplus P \to M^+_2(u,i) \oplus M^+_2(v,j) \to 0.$$

We now prove that $M^+_2(u+v, \sigma^{-1}_{u+v}(i+j+\pi))$ embeds into the projective P : if not, then the socle of $M^+_2(u+v, \sigma^{-1}_{u+v}(i+j+\pi))$, which is $L(u+v, i+j+\pi)$, embeds into one of the modules of length two, so we would have $L(u+v,i+j+\pi)=L(u+v,\sigma_{u+v}^2(i+j+\theta))=L(u+v,i+j+\theta+d)$ and hence $\pi=\theta+d$, which is impossible since π and θ are in the same range, which has at most $d-1$ elements.

Therefore $M_2^+(u+v,\sigma_{u+v}^{-1}(i+j+\pi))$ embeds into the projective P and gives $M_2^+(u+v,i+j+\pi)$ in the quotient. This means that in the quotient we have a projective, $M_2^+(u+v,i+j+\pi)$ and $M_2^+(u+v,\sigma_{u+v}(i+j+\theta))$ with θ and π in the same range. We cannot have $i+j+\pi=\sigma_{u+v}(i+j+\theta)$ (again, consider cases as in the proof of Proposition 4.17) so these modules of length two do not link up to give modules of length four. They must therefore occur as summands.

■

Proposition 4.21 *The module $M_2^-(u,i)\otimes M_2^+(v,j)$ is projective.*

Proof: If we tensor the exact sequence $0\to L(v,\sigma_v^{-1}(j))\to\Omega^{-1}(L(v,j))\to M_2^+(v,j)\to 0$ on the left by $M_2^-(u,i)$, we get the exact sequence $0\to\oplus_\pi M_2^-(u+v,\sigma_{u+v}(i+j+\pi))\to\oplus_\theta M_2^-(u+v,\sigma_{u+v}^{-1}(i+j+\theta))\oplus P\to M_2^-(u,i)\otimes M_2^+(v,j)\to 0$ with π and θ in the same range. So $M_2^-(u,i)\otimes M_2^+(v,j)$ must be a sum of modules of the form $M_2^-(u+v,\ell)$ and a projective.

A similar argument starting with the exact sequence $0\to L(u,\sigma_u(i))\to\Omega^{-1}(L(u,i))\to M_2^-(u,i)\to 0$ implies that $M_2^-(u,i)\otimes M_2^+(v,j)$ must be a sum of modules of the form $M_2^+(u+v,\ell)$ and a projective.

Combining these results implies that $M_2^-(u,i)\otimes M_2^+(v,j)$ is projective since the decomposition into indecomposable summands is unique and we can distinguish modules of the form $M_2^+(w,\ell)$ and modules of the form $M_2^-(w,\ell)$ by viewing them as $\Lambda_{n,d}$-modules and checking whether as such they are projective or not.

■

Theorem 4.22 *If ℓ and t are two positive integers, then $M_{2\ell}^+(u,i)\otimes M_{2t}^-(v,j)$ is projective, and we have the decompositions:*

$$M_{2\ell}^+(u,i)\otimes M_{2t}^+(v,j)\cong\bigoplus_{\theta\in\mathfrak{I}}\bigoplus_{p=0}^{2\ell-1}\bigoplus_{r=0}^{t-1}M_2^+(u+v,\sigma_{u+v}^{p+2r}(i+j+\theta))\oplus\text{projective}$$

and

$$M_{2\ell}^-(u,i)\otimes M_{2t}^-(v,j)\cong\bigoplus_{\theta\in\mathfrak{I}}\bigoplus_{p=0}^{2\ell-1}\bigoplus_{r=0}^{t-1}M_2^-(u+v,\sigma_{u+v}^{-(p+2r)}(i+j+\theta))\oplus\text{projective}$$

Proof: We work by induction on ℓ and t, using Proposition 3.2 (ii), since we know that modules of even length are not splitting trace modules (Proposition 3.6), so we will obtain split exact sequences.

Let us consider for instance $M_{2\ell}^+(u,i)\otimes M_{2t}^+(v,j)$. We first set $t=1$ and consider the Auslander-Reiten sequence for $M_{2\ell}^+(u,i+d)$. An induction on ℓ gives $M_{2\ell+2}^+(u,i+d)\otimes M_2^+(v,j)$ and hence $M_{2\ell}^+(u,i+d)\otimes M_2^+(v,j)$ for all ℓ.

We can then do an induction on t, considering the Auslander-Reiten sequence for $M_{2t}^+(v,j+d)$, to get the result. ■

4.5 Tensor products of band modules with modules of odd length

Consider the tensor product $C_\lambda^{1+}(u,i)\otimes L(v,j)$. Using a method similar to the proof of Proposition 4.17, we can see that this tensor product decomposes as $\bigoplus_\theta C_{\mu_\theta}^{1+}(u,i)$. We must then determine the parameters μ_θ. We will describe the method on an example, which will give an algorithm to determine these parameters for any given example, and illustrate the complexity of the general case.

Example **4.23** Let us consider the tensor product $C_\lambda^{1+}(1,5)\otimes L(0,2)$ when $n=d=6$. As we have said above, we know that this tensor product decomposes as $C_{\mu_0}^{1+}(1,1)\oplus C_{\mu_1}^{1+}(1,2)\oplus L(1,3)$ for some μ_0 and μ_1 to be determined.

The blocks involved are

$$\begin{tikzcd} L(1,5) \arrow[r, bend left=50, "b_0=\gamma^2"] \arrow[r, bend left=15, "\overline{b}_1=X^4"] & L(1,1) \arrow[l, bend left=15, "b_1=\gamma^4"] \arrow[l, bend left=50, "\overline{b}_0=12X^2"] \end{tikzcd} \quad \text{and} \quad \begin{tikzcd} L(1,2) \arrow[r, bend left=50, "b'_0=\gamma^2"] \arrow[r, bend left=15, "\overline{b}'_1=X^4"] & L(1,4). \arrow[l, bend left=15, "b'_1=\gamma^4"] \arrow[l, bend left=50, "\overline{b}'_0=12X^2"] \end{tikzcd}$$

Determine μ_0 : Recall that, for any $\lambda \in k$, the module $C_\lambda^{1+}(u,i)$ is characterised by the following pullback diagram:

$$\begin{tikzcd} 0 \arrow[r] & L(u,\sigma_u(i)) \arrow[r] \arrow[d, equal] & C_\lambda^{1+}(u,i) \arrow[r] \arrow[d] & L(u,i) \arrow[r] \arrow[d, "\varphi_\lambda"] & 0 \\ 0 \arrow[r] & L(u,\sigma_u(i)) \arrow[r] & P(u,\sigma_u(i)) \arrow[r] & \Omega^{-1}(L(u,\sigma_u(i))) \arrow[r] & 0 \end{tikzcd}$$

where the homomorphism φ_λ is determined by $\varphi_\lambda(\tilde{H}_{u,i}) = \lambda\overline{b}_0 D_{u,\sigma_u(i)} + b_1 D_{u,\sigma_u(i)}$.

We first find Z_0 in $L(1,5)\otimes L(0,2)$ corresponding to $\tilde{H}_{11}$ in $L(1,1)$ (up to scalar multiples): this satisfies $e_1 Z_0 = Z_0$ so we must have $Z_0 = \tilde{H}_{15}\otimes\tilde{H}_{02}$, and we check that $XZ_0 = 0$.

We then apply $\varphi_\lambda\otimes\mathrm{id}$ to Z_0 : this is equal to (x_0, y_0) with $x_0 = \lambda(\overline{b}_0 D_{11})\otimes\tilde{H}_{02} = 12\lambda\tilde{D}_{11}\otimes\tilde{H}_{02}$ and $y_0 = (b_1 D_{11})\otimes\tilde{H}_{02} = -12H_{11}\otimes\tilde{H}_{02}$.

We now need to lift x_0 and y_0 to $P(1,1)\otimes L(0,2)$:

- Since $Xx_0 = 0$, we know that x_0 corresponds to $\tilde{D}_{15}$, so we do not change x_0.
- We look for $\bar{y}_0 = -12H_{11}\otimes\tilde{H}_{02} + \alpha XH_{11}\otimes a\tilde{H}_{02} + \beta X^2 H_{11}\otimes\gamma^2\tilde{H}_{02}$. The element $\bar{y}_0$ is characterised by $X\bar{y}_0 \neq 0$ and $aX\bar{y}_0 = 0$ (it corresponds to H_{15}). This gives the equations $-12q^2 + \alpha(2q+q^2) + 2\beta(1+q) = 0$ and $2(q^2-1)\beta - \alpha = 0$, so finally

$$\bar{y}_0 = -12H_{11}\otimes\tilde{H}_{02} + 4(1+q)XH_{11}\otimes a\tilde{H}_{02} - 2qX^2 H_{11}\otimes\gamma^2\tilde{H}_{02}.$$

We know that $\varphi_{\mu_0}(\tilde{H}_{11}) = (\mu_0\bar{b}_1 D_{15}, b_0 D_{15})$ corresponds to $(x_0, \bar{y}_0)$, so $b_1 x_0 = \mu_0\bar{b}_0\bar{y}_0$.

Calculating $b_1 x_0$ gives $12\lambda[6X^2 H_{11}\otimes\tilde{H}_{02} - 2(q+q^2)X^3 H_{11}\otimes a\tilde{H}_{02} - 2\tilde{H}_{11}\otimes\gamma^2\otimes\tilde{H}_{02}]$, and calculating $\bar{b}_0\bar{y}_0$ gives $12[-6X^2 H_{11}\otimes\tilde{H}_{02} + 2(q+q^2)X^3 H_{11}\otimes a\tilde{H}_{02} + 2\tilde{H}_{11}\otimes\gamma^2\otimes\tilde{H}_{02}]$, so finally $\mu_0 = -\lambda$.

Determine μ_1 : We proceed in the same way: $\tilde{H}_{12}$ corresponds to $Z_1 = (1+q)a\tilde{H}_{15}\otimes\tilde{H}_{02} + \tilde{H}_{15}\otimes a\tilde{H}_{02}$ in $L(1,5)\otimes L(0,2)$, and applying $\varphi_\lambda\otimes\mathrm{id}$ gives (x_1, y_1) where $x_1 = 12\lambda[(q+q^2)XD_{11}\otimes\tilde{H}_{02} + \tilde{D}_{11}\otimes a\tilde{H}_{02}]$ and $y_1 = 12[-(q+q^2)F_{11}\otimes\tilde{H}_{02} - H_{11}\otimes a\tilde{H}_{02}]$ (note that $\varphi_\lambda(a\tilde{H}_{15}) = a\varphi_\lambda(\tilde{H}_{15})$). Lifting to $P(1,1)\otimes L(0,2)$ does not change x_1, and y_1 becomes $\bar{y}_1 = 12[-(q+q^2)F_{11}\otimes\tilde{H}_{02} - H_{11}\otimes a\tilde{H}_{02} + XH_{11}\otimes\gamma^2\tilde{H}_{02}]$.

Then, since $\varphi_{\mu_1}(\tilde{H}_{12}) = (\mu_1\bar{b}'_0 D_{14}, b'_1 D_{14})$ corresponds to a scalar multiple of $(x_1, \bar{y}_1)$, we have $b'_0 x_1 = \mu_1\bar{b}'_1\bar{y}_1$, and calculating each of the terms in this identity yields $\mu_1 = \lambda$.

Therefore $C_\lambda^{1+}(u,i)\otimes L(v,j) = C_{-\lambda}^{1+}(1,1)\oplus C_\lambda^{1+}(1,2)\oplus L(1,3)$.

Using the method described in this example, we can calculate the tensor product of a band module of minimal length $\frac{2n}{d}$ with a simple module, and hence using Auslander-Reiten sequences and the fact that modules of odd length are syzygies of simple modules, as in the previous sections, we can determine the tensor product of any band module with a module of odd length.

4.6 Tensor products of band modules with other modules of even length

Here again, we will describe the method on an example:

Example **4.24** We shall consider the tensor product $C^{1+}_{\lambda}(1,5) \otimes C^{1+}_{\mu}(0,2)$ with $n = d = 6$. Since the modules we are tensoring are periodic, their summands must also be periodic or projective, and hence of even length. From Example 4.23 and a similar calculation which gives $C^{1+}_{\lambda}(1,5) \otimes L(0,5) = C^{1+}_{-\lambda}(1,4) \oplus C^{1+}_{\lambda}(1,5) \oplus L(1,0)$, we see that the non-projective composition factors of $C^{1+}_{\lambda}(1,5) \otimes C^{1+}_{\mu}(0,2)$ are $L(1,1)$, $L(1,1)$, $L(1,5)$, $L(1,5)$, $L(1,2)$, $L(1,2)$, $L(1,4)$ and $L(1,4)$. Note that the first four composition factors and the last four composition factors must be in separate summands of $C^{1+}_{\lambda}(1,5) \otimes C^{1+}_{\mu}(0,2)$, since $\sigma_1(1) = 5$ and $\sigma_1(5) = 1$, neither of which is equal to 2 or 4. Call these summands M and N.

Determine the top of $C^{1+}_{\lambda}(1,5) \otimes C^{1+}_{\mu}(0,2)$: We shall use here the following result from [GMS, Propositions 1.1 and 1.2]:

$$\operatorname{Hom}(A \otimes B, C) \cong \operatorname{Hom}(A, B^* \otimes C)$$

for any $\mathcal{D}(\Lambda_{n,d})$-modules A, B and C, where B^* is the k-dual of B. Moreover, we can see that $C^{1+}_{\lambda}(u,i)^* \cong C^{1+}_{\lambda}(u,\sigma_u(i))$ (note that it is easy to see, using Proposition 2.20, that $L(u,i)^* \cong L(1-u,\sigma_u(i))$).

The module $L(1,1)$ is in the top of $C^{1+}_{\lambda}(1,5) \otimes C^{1+}_{\mu}(0,2)$ if, and only if, $\operatorname{Hom}(C^{1+}_{\lambda}(1,5) \otimes C^{1+}_{\mu}(0,2), L(1,1))$ is nonzero. We have

$$\begin{aligned}
\operatorname{Hom}(C^{1+}_{\lambda}(1,5) \otimes C^{1+}_{\mu}(0,2), L(1,1)) &\cong \operatorname{Hom}(C^{1+}_{\lambda}(1,5), C^{1+}_{\mu}(0,5) \otimes L(1,1)) \\
&\cong \operatorname{Hom}(C^{1+}_{\lambda}(1,5), C^{1+}_{\nu_1}(1,1)) \oplus \operatorname{Hom}(C^{1+}_{\lambda}(1,5), C^{1+}_{\nu_2}(1,2)) \\
&\quad \oplus \operatorname{Hom}(C^{1+}_{\lambda}(1,5), L(1,3)) \\
&\cong k \oplus 0 \oplus 0 = k.
\end{aligned}$$

Hence $L(1,1)$ occurs once in the top.

Similarly, $L(1,2)$ occurs once in the top. Now

$$\begin{aligned}
\operatorname{Hom}(C^{1+}_{\lambda}(1,5) \otimes C^{1+}_{\mu}(0,2), L(1,5)) &\cong \operatorname{Hom}(C^{1+}_{\lambda}(1,5), C^{1+}_{\mu}(0,5) \otimes L(1,5)) \\
&\cong \operatorname{Hom}(C^{1+}_{\lambda}(1,5), C^{1+}_{\nu_3}(1,4)) \oplus \operatorname{Hom}(C^{1+}_{\lambda}(1,5), C^{1+}_{\nu_4}(1,5)) \\
&\quad \oplus \operatorname{Hom}(C^{1+}_{\lambda}(1,5), L(1,0)) \\
&\cong \operatorname{Hom}(C^{1+}_{\lambda}(1,5), C^{1+}_{\nu_4}(1,5)) \\
&= \begin{cases} k & \text{if } \nu_4 = \lambda \\ 0 & \text{otherwise,} \end{cases}
\end{aligned}$$

since any homomorphism must be an isomorphism. So we need to determine ν_4. Proceeding as in Example 4.23 gives $\nu_4 = -\frac{1+q}{2}\mu$. Hence $L(1,5)$ occurs in the top if, and only if, $\lambda = -\frac{1+q}{2}\mu$.

Similarly, $L(1,4)$ occurs in the top if, and only if, $\lambda = \frac{1+q}{2}\mu$.

Determine the socle of $C^{1+}_{\lambda}(1,5) \otimes C^{1+}_{\mu}(0,2)$: This is similar, considering $\operatorname{Hom}(L(u,i), C^{1+}_{\lambda}(1,5) \otimes C^{1+}_{\mu}(0,2))$ for each of the simples $L(u,i)$ involved. We find that $L(1,1)$ and $L(1,2)$ both occur once in the socle, $L(1,5)$ occurs in the socle if, and only if, $\lambda = -\frac{1+q}{2}\mu$, and $L(1,4)$ occurs in the socle if, and only if, $\lambda = \frac{1+q}{2}\mu$.

First case: $\lambda \neq \pm\frac{1+q}{2}\mu$: Then $L(1,4)$ and $L(1,5)$ do not occur in the top or the socle of $C^{1+}_{\lambda}(1,5) \otimes C^{1+}_{\mu}(0,2)$. Therefore the two summands we considered above each have a simple top and a simple socle, so $C^{1+}_{\lambda}(1,5) \otimes C^{1+}_{\mu}(0,2) = P(1,1) \oplus P(1,2) \oplus$ projective is a projective module.

Second case: $\lambda = \frac{1+q}{2}\mu$: Then the summand M is as in the first case and must be $P(1,1)$, but the summand N differs: $L(1,2)$ and $L(1,4)$ both occur once in the top and once in the socle, and since the summands must have even length, N must be $C^{1+}_{\nu_5}(1,2) \oplus C^{1+}_{\nu_6}(1,4)$.

In order to determine ν_5 and ν_6, consider the following exact sequence:

$$0 \to L(0,5) \to C^{1+}_{\mu}(0,2) \to L(0,2) \to 0.$$

Tensoring with $C^{1+}_{\lambda}(1,5)$ gives the exact sequence:

$$0 \to C^{1+}_{-\lambda}(1,4) \oplus C^{1+}_{\lambda}(1,5) \oplus L(1,0) \to C^{1+}_{\lambda}(1,5) \otimes C^{1+}_{\mu}(0,2) \to C^{1+}_{-\lambda}(1,1) \oplus C^{1+}_{\lambda}(1,2) \oplus L(1,3) \to 0,$$

so we know, by considering the composition factors of $C^{1+}_{-\lambda}(1,4)$ and of $C^{1+}_{\lambda}(1,5) \otimes C^{1+}_{\mu}(0,2)$, that $C^{1+}_{-\lambda}(1,4)$ embeds into $C^{1+}_{\nu_5}(1,2) \oplus C^{1+}_{\nu_6}(1,4)$. So we have an embedding $\iota : C^{1+}_{-\lambda}(1,4) \to C^{1+}_{\nu_5}(1,2) \oplus C^{1+}_{\nu_6}(1,4)$ and a projection $\pi : C^{1+}_{\nu_5}(1,2) \oplus C^{1+}_{\nu_6}(1,4) \to C^{1+}_{\nu_6}(1,4)$. The composition $\pi\iota$ is non-zero on the socle $L(1,2)$, hence injective, so it must be an isomorphism. Therefore we have $\nu_6 = -\lambda$. A similar argument, using the fact that $C^{1+}_{\lambda}(1,2)$ is a quotient of $C^{1+}_{\lambda}(1,5) \otimes C^{1+}_{\mu}(0,2)$, shows that $\nu_5 = \lambda$.

Finally $C^{1+}_{\lambda}(1,5) \otimes C^{1+}_{\mu}(0,2) = C^{1+}_{\lambda}(1,2) \oplus C^{1+}_{-\lambda}(1,4) \oplus$ projective.

Third case: $\lambda = -\dfrac{1+q}{2}\mu$: This is similar to the second case, and we get $C^{1+}_{\lambda}(1,5) \otimes C^{1+}_{\mu}(0,2) = C^{1+}_{-\lambda}(1,1) \oplus C^{1+}_{\lambda}(1,5) \oplus$ projective.

Other situations (such as $d \neq n$ or tensor products of different types of band modules of minimal length) are treated similarly. Moreover, the case of the tensor product of a band module with a string module of even length can also be treated in this way, setting the parameter to zero for the string module. Finally, the general case (that is, when the lengths are not minimal) can be deduced by induction, using Auslander-Reiten sequences.

5 Interpretation of simple and projective $\mathcal{D}(\Lambda_{n,d})$-modules as Hopf bimodules over $\Lambda_{n,d}$

In this section, we describe some Hopf bimodules over $\Lambda_{n,d}$: given any finite-dimensional Hopf algebra H, there is an equivalence of categories between modules over the Drinfel'd double $\mathcal{D}(H)$ and Hopf bimodules over H (see [Kl, M, R] for instance). Recall:

Definition 5.1 *Let H be a Hopf algebra. A* **Hopf bimodule** *over H is an H-bimodule M which is an H-bicomodule such that the comodule structure maps $M \to H \otimes M$ and $M \to M \otimes H$ are H-bimodule homomorphisms ($H \otimes M$ and $M \otimes H$ are endowed with diagonal H-bimodule structures). We use Sweedler's notation for the comultiplication of H, so $\Delta(h) = h^{(1)} \otimes h^{(2)}$.*

The functor from $\mathcal{D}(H)$-modules to Hopf bimodules over H can be described as follows: let V be a module over $\mathcal{D}(H)$. Consider the vector space $M := H \otimes V$; as a right H-module, it is free (of rank $\dim V$), that is, for any $h, a \in H$ and $v \in V$, we have $(h \otimes v) \cdot a = ha \otimes v$. As a left H-comodule, it is free, that is, $h \otimes v \mapsto h^{(1)} \otimes (h^{(2)} \otimes v)$. The left module structure on M is diagonal, that is, $a \cdot (h \otimes v) = a^{(1)}h \otimes a^{(2)}v$.

Finally, to describe the right comodule structure, we need to fix a basis $\{b_i\}$ of H and $\{b_i^*\}$ its dual basis; then $\delta_R(h \otimes v) = \sum_i h^{(1)} \otimes b_i^* \cdot v \otimes b_i \cdot h^{(2)}$. Note that V is a left H^*-module and a left H-module (restricting the action of $\mathcal{D}(H)$ to H^{*cop} and to H).

If we take $H = \Lambda_{n,d}$, we have a basis given by $\{q^{-im}\frac{1}{m!_q}\gamma_i^m \mid i \in \mathbb{Z}_n, 0 \leqslant m \leqslant d-1\}$. We refer the reader to the notation and correspondence in Section 2 (page 3) in order to see that the dual basis is $\{X^m G^i \mid i \in \mathbb{Z}_n, 0 \leqslant m \leqslant d-1\}$. Hence, if V is a module over $\mathcal{D}(\Lambda_{n,d})$, we have

$$\delta_R(h \otimes v) = \sum_{\substack{i \in \mathbb{Z}_n \\ 0 \leqslant m \leqslant d-1}} (h^{(1)} \otimes X^m G^i v) \otimes q^{-im} \frac{1}{m!_q} \gamma_i^m h^{(2)}.$$

Therefore, if V is a module over $\mathcal{D}(\Lambda_{n,d})$, the corresponding Hopf bimodule over $\Lambda_{n,d}$ is free as a right module and left comodule, so as such it is isomorphic to $\Lambda_{n,d}^{\dim V}$, and we now need to describe the left module and right comodule structures on $\Lambda_{n,d}^{\dim V}$ using the definitions above.

5.1 Hopf bimodule corresponding to $L(u, j)$

Let N be the dimension of $L(u, j)$. Let us first describe the left $\Lambda_{n,d}$-module structure of $\Lambda_{n,d} \otimes L(u, j)$.

Define $\tau_k^{(p)}(\gamma_i^m) = \binom{m}{p}_q q^{(m-p)k}\gamma_{i-k}^{m-p}$. Then, using the definitions above, we can easily see that

$$\gamma_i^m \cdot (h \otimes \gamma_j^t \tilde{H}_{u,j}) = \sum_{p=0}^{m} \tau_{j+t}^{(p)}(\gamma_i^m) h \otimes \gamma_j^{t+p} \tilde{H}_{u,j}.$$

Therefore, as a left $\Lambda_{n,d}$-module, $\Lambda_{n,d} \otimes L(u, j)$ is isomorphic to ${}_{T_j(N)}(\Lambda_{n,d}^N)$, that is, the left module whose action is described by

$$\Lambda_{n,d} \otimes \Lambda_{n,d}^N \overset{T_j(N)\otimes 1}{\longrightarrow} \mathcal{M}_N(\Lambda_{n,d}) \otimes \Lambda_{n,d}^N \overset{\text{matrix multiplication}}{\longrightarrow} \Lambda_{n,d}^N$$

where $T_j(N) : \Lambda_{n,d} \to \mathcal{M}_N(\Lambda_{n,d})$ is defined by $T_j(N) = \begin{pmatrix} \tau_j^{(0)} & \tau_j^{(1)} & \cdots & \tau_j^{(N-1)} \\ & \tau_{j+1}^{(0)} & & \vdots \\ & 0 & \ddots & \vdots \\ & & & \tau_{j+N-1}^{(0)} \end{pmatrix}$. Note that if $\dim L(u, j) = 1$, this describes a twisted module ${}_{\tau}(\Lambda_{n,d})$, where τ is an automorphism of $\Lambda_{n,d}$.

We proceed in a similar way for the right coaction. Let $R_j(N)$ be an element in $\mathcal{M}_N(\Lambda_{n,d})$. Define a right coaction on $\Lambda_{n,d}^N$ as follows:

$$\Lambda_{n,d}^N \xrightarrow{\Delta} (\Lambda_{n,d} \otimes \Lambda_{n,d})^N \overset{(1\otimes R_j(N))\cdot}{\longrightarrow} (\Lambda_{n,d} \otimes \Lambda_{n,d})^N \cong \Lambda_{n,d}^N \otimes \Lambda_{n,d}.$$

Setting $R_j(N) = \begin{pmatrix} \rho_{j+N-1}^{(0)} & & & \\ \rho_{j+N-2}^{(1)} & \rho_{j+N-2}^{(0)} & 0 & \\ \vdots & & \ddots & \\ \rho_j^{(d+N-1)} & & & \rho_j^{(0)} \end{pmatrix}$ with $\rho_k^{(0)} = \sum_{i\in\mathbb{Z}_n} q^{-ik} e_i$ and

$$\rho_k^{(m)} = \binom{-k+N+j-1-m}{m}_q \left(\prod_{p=-k+j-m}^{-k+j-1} (1 - q^{p+m}) \right) \sum_{i\in\mathbb{Z}_n} q^{-ik}\gamma_i^m \text{ for } m \geqslant 1$$

finally gives:

Proposition 5.2 *The Hopf bimodule corresponding to the $\mathcal{D}(\Lambda_{n,d})$-module $L(u, j)$ is*

$${}^{1}_{T_j(N)}[(\Lambda_{n,d})^{\dim L(u,j)}]_1^{R_j(N)}.$$

Note that when $\dim L(u, j) = 1$, we have ${}^{1}_{T_j(1)}[(\Lambda_{n,d})^{\dim L(u,j)}]_1^{R_j(1)}$, that is, the usual Hopf bimodule $\Lambda_{n,d}$ twisted on the left by the algebra automorphism $T_j(1)$ and on the right by the coalgebra automorphism $R_j(1)\cdot$ (in this case $R_j(1) = \rho_j^{(0)}$ is a grouplike element). In particular, the trivial module $L(0, 0)$ gives the usual Hopf bimodule $\Lambda_{n,d}$.

5.2 Hopf bimodule corresponding to $P(u, j)$

We assume here that $P(u, j)$ is not simple, that is, $N = \dim L(u, j) \neq d$. We then have a basis for $P(u, j)$ given by $\{A_\ell = \gamma_j^\ell D_{uj}, B_\ell = \gamma_{j+N}^\ell X_{uj}^{d-N} \mid 0 \leqslant \ell \leqslant d-1\}$. It is easy to work out the actions of G and of paths on these basis elements, and using Lemma 2.17, we can also describe the action of X. We then obtain:

Proposition 5.3 *The Hopf bimodule corresponding to the* $\mathcal{D}(\Lambda_{n,d})$*-module* $P(u,j)$ *is*

$$ {}^{1}_{U_j}[(\Lambda_{n,d})^{2d}]^{V_j}_{1}, $$

where $U_j = \begin{pmatrix} T_j(d) & 0 \\ 0 & T_{j+N}(d) \end{pmatrix}$ *and* $V_j = \begin{pmatrix} R_j(d) & 0 \\ \Omega_j & \Pi_j \end{pmatrix}$ *for some matrices* Ω_j *and* Π_j *which can be explicitly calculated, with* Π_j *lower triangular.*

A Appendix: Proof of the classification of the indecomposable $\mathcal{D}(\Lambda_{n,d})$-modules

We wish to describe all the indecomposable modules over $\mathcal{D}(\Lambda_{n,d})$ up to isomorphism. We may restrict our study to a nonsemisimple block $\mathbb{B}$ of $\mathcal{D}(\Lambda_{n,d})$. To simplify notation, we set $m := \frac{2n}{d}$ to be the number of simple modules over $\mathbb{B}$, which we denote by $S_p := L(u, \sigma_u^p(i))$ with $p \in \mathbb{Z}_m$. Let ϵ_p be the corresponding idempotent, and P_p the projective cover of S_p.

We first describe the Loewy structure of the indecomposable modules, then describe the indecomposable modules themselves. Finally, we give a different but more elegant proof of the fact that indecomposable modules of odd length are syzygies of simple modules (this proof is valid for any m, not necessarily even).

A.1 Loewy structure of indecomposable modules

Lemma A.1 *Let* $\mathbb{B}$ *be a fixed block as above, with* m *simple modules where* $m \geqslant 2$*. Then for any indecomposable* $\mathbb{B}$*-module* M *which is not simple or projective, the radical of* M *is equal to the socle of* M*.*

Proof: All the indecomposable projective modules have Loewy length three, so an indecomposable non-projective module has Loewy length at most 2, and equal to 2 if it is not simple.

Moreover $\mathrm{soc}(M) \subseteq \mathrm{rad}(M)$ in general: indeed, if not, there would be an element x in the socle which is not in the radical. Choose x such that $\mathbb{B}x$ is simple. Take a maximal submodule $\mathfrak{M}$ of M which does not contain x; then $\mathfrak{M} \cap M = 0$ so $M = \mathfrak{M} \oplus \mathbb{B}x$ is decomposable.

Therefore, if the Loewy length of M is 2, then $\mathrm{soc}(M)$ and $\mathrm{rad}(M)$ must be equal. ■

We now describe the top and the radical of an indecomposable module.

Lemma A.2 *Suppose* M *is indecomposable and not projective or simple. One of the following must hold:*

- *(i)* $M/\mathrm{rad}(M) = \oplus_p \epsilon_{2p} M/\mathrm{rad}(M)$ *and* $\mathrm{rad}(M) = \oplus_p \epsilon_{2p+1}\mathrm{rad}(M)$*;*
- *(ii)* $M/\mathrm{rad}(M) = \oplus_p \epsilon_{2p+1} M/\mathrm{rad}(M)$ *and* $\mathrm{rad}(M) = \oplus_p \epsilon_{2p}\mathrm{rad}(M)$*.*

Proof: Choose a vector space complement C with $M = \mathrm{rad}(M) \oplus C$ such that $\epsilon_p C \subseteq C$ for all p (this is possible since the algebra generated by the idempotents ϵ_p is semi-simple). Then if α is any arrow starting at p we have $\alpha C = \alpha\epsilon_p C \subseteq \epsilon_{p+1}\mathrm{rad}(M) \oplus \epsilon_{p-1}\mathrm{rad}(M)$. But $p \pm 1$ have the same parity and m is even, so if

$$M' := [\oplus_p \epsilon_{2p} C] \oplus [\oplus_p \epsilon_{2p+1}\mathrm{rad}(M)], \quad M'' := [\oplus_p \epsilon_{2p+1} C] \oplus [\oplus_p \epsilon_{2p}\mathrm{rad}(M)]$$

then these are $\mathbb{B}$-submodules of M (use Lemma A.1) and $M = M' \oplus M''$. If $M = M'$ then we have (i) and otherwise (ii). ■

Definition A.3 *We say that the top of* M *is* **even** *if Lemma A.2(i) holds, and that the top of* M *is* **odd** *if Lemma A.2(ii) holds.*

A.2 Description of the indecomposable modules over a block $\mathbb{B}$

A.2.1 Applying Fitting's Lemma

Let M be an indecomposable module. We can assume, without loss of generality, that the top of M is even. Set $M_p = \epsilon_p M$: this is $\epsilon_p M/\mathrm{rad}(M)$ if p is even, and $\epsilon_p \mathrm{rad}(M)$ otherwise.

Then M is completely described by the linear maps given by left multiplication,

$$b_{2r} : M_{2r} \to M_{2r+1}, \quad \overline{b}_{2r-1} : M_{2r} \to M_{2r-1}$$

(the other arrows act as zero). We would like to have maps which we can compose, so we define $\beta_p := {}^t(\overline{b}_p) : M_p^* \to M_{p+1}^*$, for p odd. For any fixed basis of M_p and M_{p+1}, with respect to the dual bases, the matrix of the map β_p is the transpose of the matrix of $\overline{b}_p$. Let b and β also denote the matrices of the maps b and β.

Consider the map $\alpha_p : M_p \to M_p$, which is equal to $\beta_{p-1} b_{p-2} \dots b_{p+2} \beta_{p+1} b_p$ if p is even, and to $b_{p-1} \beta_{p-2} \dots \beta_{p+2} b_{p+1} \beta_p$ if p is odd, that is, the composition of all b's and β's, one each in the only possible order.

By Fitting's Lemma, there is some large $N > 0$ such that $U_p := \mathrm{Ker}(\alpha_p^N) = \mathrm{Ker}(\alpha_p^{N+\ell})$ and $V_p := \mathrm{Im}(\alpha_p^N) = \mathrm{Im}(\alpha_p^{N+\ell})$ for all $\ell \geq 0$, where U_p and V_p are α_p-invariant and $M_p = U_p \oplus V_p$. Moreover $\alpha_p|_{U_p}$ is nilpotent and $\alpha_p|_{V_p}$ is an isomorphism.

Set $U := \oplus_p U_p$ and $V := \oplus_p V_p$. Then $M = U \oplus V$ and moreover, both U and V are invariant under the actions of b_{2r} and β_{2r+1}. Let us check this for b_{2r} : note that $\alpha_{2r+1} b_{2r} = b_{2r} \alpha_{2r}$. So if x is in U_{2r}, then $\alpha_{2r+1}^N b_{2r}(x) = b_{2r} \alpha_{2r}^N(x) = 0$, so $b_{2r}(x)$ is in U_{2r+1}. If x is in V_{2r}, then we can write $x = \alpha_{2r}^N(z)$ and hence $b_{2r}(x) = \alpha_{2r+1}^N(z) \in V_{2r+1}$.

The decomposition $M = U \oplus V$ above is a decomposition of $\mathbb{B}$-modules: for each M_p take a basis for U_p and a basis for V_p such that the union is a basis for M_p. Then, for each t odd, the matrix of β_t is a block-diagonal matrix. Now return to $\overline{b}_t$ (instead of β_t): we take the transpose of this matrix, which has the same block form. Therefore $M = U \oplus V$ as a $\mathbb{B}$-module.

A.2.2 Canonical forms

We now assume that M is an indecomposable module. Therefore, M is equal to U or V.

First case: $M = U$. Consider the algebra A generated by the β_{2p-1} and the b_{2p}. This is a Nakayama algebra, whose quiver is cyclic, and all the paths of length at least δ where $\delta = m \cdot \max\{\text{nilpotence degree of the } \alpha_p\}$ are zero, with $\delta \geqslant \dim M$.

We know that modules over A are uniserial. Therefore there is a generator v of M over A such that $v = \epsilon_p v$, and a basis for M is given by $\{v, b_p v, \beta_{p+1} b_p v, \dots\}$ (the index p is even here, since we are in the even top case).

Returning to b's and $\overline{b}$'s, we can write the matrices for the actions of these arrows, and we see that M is a string module. If it has odd length then we recover a syzygy of a simple module, and otherwise we have a string module of type $M_2^+(u,i)$. Note that if we had started with a module with odd top, we would have obtained either a syzygy of a simple module (if the length were odd), or a string module of type $M_2^-(u,i)$ (if the length were even). See Section 3.1 for the notations.

Second case: $M = V$. We have $\mathrm{Ker}(\alpha_p^N) = 0$ for all p and $M = \bigoplus_p \mathrm{Im}(\alpha_p^n)$, so we deduce that the maps b_{2r} and the maps β_{2r-1} (and hence $\overline{b}_{2r-1}$) are invertible. Let ℓ be the dimension of the M_p (they all have the same dimension). We can choose bases for the M_p such that all the maps b_{2r} and $\overline{b}_{2r-1}$ are represented by the identity matrix, except for one of them, say b_0. Note that the bases for the M_p are determined by a choice of basis for M_1 say, and any change of basis for M_1 leads to the same change of basis for all the other M_p. Since M is indecomposable, the map b_0 must be an indecomposable linear map, so we can choose a basis for M_1 such that the matrix of b_0 is of the form $J_\ell(\lambda) = \begin{pmatrix} \lambda & 1 & \dots & \\ 0 & \lambda & 1 & \dots \\ & & \dots & \\ 0 & \dots & & \lambda \end{pmatrix}$ for some nonzero λ in k, and such that all the other matrices remain identity matrices.

We call this module $C^{\ell+}_{\lambda}(u,i)$. The same construction starting with a module whose top is odd will give us $C^{\ell-}_{\lambda}(u,i)$. Note that they have even length, equal to $m\ell$.

A.2.3 Auslander-Reiten components

We refer the reader to [ARS, V.1 and VII.1] and [E, I.7 and I.8] for information about Auslander-Reiten sequences and quivers.

(I) Indecomposable modules of odd length: For each simple module S_p we have the standard Auslander-Reiten sequence

$$0 \to \Omega(S_p) \to \mathrm{rad}(P_p)/\mathrm{soc}(P_p) \oplus P_p \to \Omega^{-1}(S_p) \to 0$$

These are the only Auslander-Reiten sequences where projectives occur. For a symmetric algebra (which is the case here, see the Introduction), the Auslander-Reiten translation is the same as Ω^2, so it follows from Section A.2.2 that any indecomposable module of odd length, say $M = \Omega^k(S_p)$ with $k \neq -1$, has Auslander-Reiten sequence

$$0 \to \Omega^{k+2}(S_p) \to \Omega^{k+1}(S_{p-1}) \oplus \Omega^{k+1}(S_{p+1}) \to M \to 0,$$

obtained by applying Ω^{k+1} to the sequence above.

This describes all the Auslander-Reiten sequences, and therefore the Auslander-Reiten component in which M occurs.

(II) String modules of even length: By explicit calculation, we can easily see that $\Omega(M^+_{2\ell}(u,i)) = M^+_{2\ell}(u,\sigma_u^{-1}(i))$. So $M^+_{2\ell}(u,i)$ has Ω-period equal to m, hence it lies in a tube of rank $\frac{m}{2} = \frac{n}{d}$.

We now determine the Auslander-Reiten sequences. Let us start with the Auslander-Reiten sequence for $M^+_2(u,i)$: it is of the form $0 \to M^+_2(u,i-d) \to M \to M^+_2(u,i) \to 0$ since the Auslander-Reiten translate is given by Ω^2. The module M must have length four, and cannot be a direct sum of modules of length two (otherwise the exact sequence would split). The composition factors of M are $L(u,\sigma_u^{-2}(i))$, $L(u,\sigma_u^{-1}(i))$, $L(u,i)$ and $L(u,\sigma_u(i))$. Moreover, $L(u,\sigma_u^{-2}(i))$ is in the socle of $M^+_2(u,i-d)$ so it must be in the socle of M, and $L(u,i)$ is in the top of $M^+_2(u,i)$ so it must be in the top of M. It then follows from the classification of the modules in Section A.2.2 that M must be $M^+_4(u,i-d)$.

Let us now determine the Auslander-Reiten sequence for $M^+_4(u,i)$. We know that the left hand term of the sequence must be $\Omega^2(M^+_4(u,i))$, and we know exactly which modules of length two occur, from the Auslander-Reiten sequences for modules of length two. Therefore we have $0 \to M^+_4(u,i-d) \to M^+_2(u,i) \oplus N \to M^+_4(u,i) \to 0$, with N indecomposable of length six. The composition factors of N are $L(u,\sigma_u^{-1}(i))$, $L(u,\sigma_u(i))$, $L(u,\sigma_u^3(i))$, $L(u,\sigma_u^{-2}(i))$ $L(u,i)$ $L(u,\sigma_u^2(i))$. The first one must be in the socle, and hence so must the next two. The last one must be in the top, and therefore so are the two remaining simple modules. Therefore $N = M^+_6(u,i)$.

The general sequences in Section 3.2**(II)** are obtained in this way by induction. Therefore, the $M^+_{2\ell}(u,\sigma_u^p(i))$ for all even p and all ℓ form precisely one tube, where the modules of length 2ℓ form the ℓ^{th} row. The $M^+_{2\ell}(u,\sigma_u^p(i))$ for all odd p and all ℓ form precisely one tube, where the modules of length 2ℓ form the ℓ^{th} row.

Similarly, there are two tubes, each of rank $\frac{n}{d}$, containing all the modules $M^-_{2\ell}(u,\sigma_u^p(i))$.

(III) Band modules: We can see that $\Omega(C^{\ell+}_{\lambda}(u,i)) \cong C^{\ell-}_{-\lambda}(u,i)$ and $\Omega(C^{\ell-}_{\mu}(u,i)) \cong C^{\ell+}_{-\mu}(u,i)$. Hence $C^{\ell+}_{\lambda}(u,i)$ has period two and lies in a tube of rank one.

We indicate briefly how to determine the Auslander-Reiten sequences given in Section 3.2**(III)**: the Auslander-Reiten sequence for $C^{1+}_{\lambda}(u,i)$ is of the form $0 \to C^{1+}_{\lambda}(u,i) \to M \xrightarrow{\pi} C^{1+}_{\lambda}(u,i) \to 0$ (recall that the Auslander-Reiten translate is given by Ω^2 and that these modules are periodic of period two). The module M must be indecomposable of length $\frac{4n}{d}$, its top must be $\bigoplus_p L(u,\sigma_u^{2p}(u,i))^2$ and its socle is $\bigoplus_p L(u,\sigma_u^{2p+1}(u,i))^2$ (as vector spaces), so N is of the form $C^{2+}_{\mu}(u,i)$. We need to determine μ. Let $\{v_1,v_2\}$ be a basis for

$L(u,i)^2$ such that the action of b_0 is given by the matrix $\begin{pmatrix} \mu & 1 \\ 0 & \mu \end{pmatrix}$. The vector v_1 generates a submodule so must come from the left hand term in the sequence. Hence $\pi(v_1) = 0$, and $\pi(v_2) \neq 0$ is a basis for $L(u,i)$ in $C_\mu^{2,+}(u,i)$. We then have $\pi(b_0 v_2) = \pi(v_1 + \mu v_2) = \mu\pi(v_2)$, and $\pi(b_0 v_2) = b_0 \pi(v_2) = \lambda\pi(v_2)$ since the action of b_0 on $L(u,i)$ is given by multiplication by λ. Hence $\mu = \lambda$. An induction gives the expected sequences, and therefore, for each λ, the modules $C_\lambda^{\ell+}(u,i)$ form one tube, with $C_\lambda^{\ell+}(u,i)$ in row ℓ. Similarly, the $C_\lambda^{\ell-}(u,i)$ when ℓ varies form one tube, with $C_\lambda^{\ell-}(u,i)$ in row ℓ.

A.3 Indecomposable modules of odd length: another proof

We work over a block $\mathbb{B}$ with m simple modules, where m is any integer at least equal to 2.

Lemma A.4 *Suppose M is indecomposable in $\mathbb{B}$ of odd length. Then some syzygy $\Omega^k(M)$ for some $k \in \mathbb{Z}$ is simple.*

Proof: Since M has odd length, $\mathrm{soc}(M)$ and $M/\mathrm{soc}(M)(= \mathrm{top}(M))$ have different length, say $\mathrm{soc}(M)$ has larger length. Consider a projective cover

$$0 \to \Omega(M) \to P \to M \to 0.$$

We know from the structure of projective modules that $\mathrm{top}(P) \cong \mathrm{soc}(P)$, so $\mathrm{top}(M) \cong \mathrm{soc}(\Omega(M))$.

If $\mathrm{top}(M)$ has length t, then P has length $4t$ (since $\mathrm{top}(P) = \mathrm{top}(M)$). It follows that $\mathrm{soc}(\Omega(M))$ has length t and $\mathrm{top}(\Omega(M))$ has length $t + x$ if $\Omega(M)$ is not simple.

Therefore the length of the module $\Omega(M)$ is strictly smaller than that of M, and $\mathrm{top}(\Omega(M))$ has more composition factors than $\mathrm{soc}(\Omega(M))$. If $\Omega(M)$ is simple then the proof is finished. Otherwise we can repeat the argument, and since the lengths are positive integers, the process stops after a finite number of steps, that is, some $\Omega^k(M)$ is simple.

If the length of $\mathrm{soc}(M)$ is smaller than the length of $\mathrm{top}(M)$, a similar argument using injective envelopes gives the result. ■

Remark **A.5** It follows from the proof above that $|\mathrm{length}(\mathrm{top}(M)) - \mathrm{length}(\mathrm{soc}(M))| = 1$ for any indecomposable module M of odd length.

References

[AC] M. Auslander and J. Carlson, Almost-split Sequences and Groups Rings, *J. ALgebra* **103** (1986), pp 122-140.

[ARS] M. Auslander, I. Reiten and S.O. Smalø, Representation Theory of Artin Algebras, *Cambridge Studies in Advanced Mathematics* **36**, *Cambridge University Press* (1995).

[B] D. Benson, Representations and Cohomology I, *Cambridge Studies in Advanced Mathematics* **30**, *Cambridge University Press* (1991).

[CP] V. Chari and A. Pressley, A Guide to Quantum Groups, *Cambridge University Press* (1994).

[Ch] H-X. Chen, Irreducible Representations of a Class of Quantum Doubles, *J. Algebra* **225** (2000), pp 391–409.

[CHZ] X-W. Chen, H-L. Huang, Y. Ye and P. Zhang, Monomial Hopf algebras, *J. Algebra* **275** (2004), pp 212-232.
****

[Ci] C. Cibils, A Quiver Quantum Group, *Comm. Math. Phys.* **157** (1993), no. 3, pp 459–477.

[E] K. Erdmann, Blocks of Tame Representation Type and Related Algebras, *Springer Lecture Notes in Mathematics* **1428**, *Springer-Verlag* (1990).

[F] R. Farnsteiner, On Frobenius Extensions Defined by Hopf Algebras, *J. Algebra* **166** (1994), pp 130–141.

[FS] R. Farnsteiner and A. Skowroński, The Tame Infinitesimal Groups of Odd Characteristic, *preprint.*

[G] E.L. Green, Introduction to Koszul Algebras, in *Representation Theory and Algebraic Geometry, London Math. Soc. Lecture Notes series* **238**, *CUP* (1997), pp 45-62.

[GMS] E.L. Green, E.N. Marcos and Ø. Solberg, Representations and Almost Split Sequences for Hopf Algebras, *Representation theory of algebras (Cocoyoc, 1994), CMS Conf. Proc.* **18**, *Amer. Math. Soc., Providence, RI* (1996), pp 237–245.

[GM] E.L. Green and R. Martínez-Villa, Koszul and Yoneda algebras, *Representation theory of algebras (Cocoyoc, 1994), CMS Conf. Proc.* **18**, *Amer. Math. Soc., Providence, RI, (1996)* pp 247–297.

[H] B. Huber, Auslander-Reiten-Folgen über Smashprodukten, *Diplomarbeit im Fach Mathematik, Universität München* (2001).

[Kl] C. Kassel, Quantum Groups, *Graduate Texts in Mathematics*, **155**, *Springer-Verlag, New York* (1995).

[Ka] S. Kawata, On Auslander-Reiten components of certain group modules, *Osaka J. Math.* **30** no.2 (1993), pp 137–157.

[M] S. Montgomery, Hopf Algebras and Their Actions on Rings, *CBMS Regional Conference Series in Mathematics*, **82** *AMS* (1993).

[NZ] W.D. Nichols and M.B. Zoeller, A Hopf Algebra Freeness Theorem, *Amer. J. Math* **111** (1989), pp 381–385.

[P] M.K. Patra, On the Structure of Nonsemisimple Hopf Algebras, *J. Phys. A. Math. Gen.* **32** (1999), pp 159–166.

[Rd] D.E. Radford, Minimal Quasitriangular Hopf Algebras, *J. Algebra* **157** (1993) no.2, pp 285–315.

[Rd2] D.E. Radford, On Oriented Quantum Algebras Derived from Representations of the Quantum Double of a Finite-dimensional Hopf Algebra, *J. Algebra* **270** (2003), pp 670–695.

[Rd3] D.E. Radford, Finite-dimensional Simple-pointed Hopf Algebras, *J. Algebra* **211** (1999), pp 686–710.

[R] Rosso, M., Groupes Quantiques et Algèbres de Battage Quantiques, *C. R. Acad. Sci. Paris Sér. I* **320** (1995), pp 145–148.

[S] R. Suter, Modules for $\mathfrak{U}_q(\mathfrak{sl}_2)$, *Comm. Math. Phys.* **163** (1994), pp 359–393.

[W] S. Witherspoon, The Representation Ring of the Quantum Double of a Finite Group, *J. Algebra* **179** (1996), pp 305–329.

[X] J. Xiao, Finite-dimensional Representations of $U_t(sl(2))$ at Roots of Unity, *Can. J. Math.* **49** (1997), no. 4, pp 772-787.

Karin Erdmann
Mathematical Institute,
24-29 St. Giles,
Oxford OX1 3LB,
United Kingdom.
E-mail: erdmann@maths.ox.ac.uk

Edward L. Green
Department of Mathematics,
Virginia Polytechnic Institute and State University,
Blacksburg, VA 24061-0123

USA.
E-mail: green@math.vt.edu

Nicole Snashall
Department of Mathematics,
University of Leicester,
Leicester LE1 7RH,
United Kingdom.
E-mail: N.Snashall@mcs.le.ac.uk

Rachel Taillefer
Laboratoire d'Arithmétique et d'Algèbre,
Faculté des Sciences et Techniques,
23 Rue Docteur Paul Michelon,
42023 Saint-Etienne Cedex 2,
France.
E-mail: rachel.taillefer@univ-st-etienne.fr
Telephone:+33 (0)4 77 48 15 33
Fax:+33 (0)4 77 48 51 33.